



BRITISH LIVER TRUST – PUBLIC POLLING MAY 2017

Methodology: ComRes interviewed 2,024 GB adults online between 24th and 25th May 2017. Data were weighted to be demographically representative of all GB adults aged 18+ by age, gender, region and socio-economic grade. ComRes is a member of the British Polling Council and abides by its rules.

All press releases or other publications must be checked with ComRes before use. ComRes requires 48 hours to check a press release unless otherwise agreed.

For information about commissioning research please contact simon.carter@comresglobal.com or call +44 (0)20 7871 8660.

To register for Pollwatch, featuring commentary and insight from the ComRes team, please email: pollwatch@comresglobal.com

REPUTATION | COMMUNICATIONS | PUBLIC POLICY

Four Millbank | London | SW1P 3JA | T. +44 (0)20 7871 8660

Rond-Point Schuman / Schumanplein 6 | Box 5 | 1040 Bruxelles | T. +32 (0)2 234 63 82

51/F Raffles City | No.268 Xi Zang Middle Road | HuangPu District Shanghai | 200001 China | T. +86 (0)21 2312 7674

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 1

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Summary table

	<u>Liver disease</u>	<u>Kidney disease</u>	<u>Breast cancer</u>	<u>Heart disease</u>	<u>Diabetes</u>	<u>Stroke</u>	<u>Road traffic accident</u>
Unweighted base	2024	2024	2024	2024	2024	2024	2024
Weighted base	2024	2024	2024	2024	2024	2024	2024
1st - would cause largest no. of deaths	51 3%	34 2%	183 9%	1185 59%	148 7%	125 6%	298 15%
2nd largest	176 9%	79 4%	356 18%	379 19%	291 14%	589 29%	154 8%
3rd largest	278 14%	187 9%	402 20%	171 8%	334 16%	428 21%	225 11%
4th largest	398 20%	317 16%	371 18%	91 4%	270 13%	351 17%	220 11%
5th largest	466 23%	460 23%	291 14%	75 4%	299 15%	222 11%	205 10%
6th largest	430 21%	526 26%	265 13%	62 3%	263 13%	219 11%	252 12%
7th - would cause the smallest no. of deaths	218 11%	415 21%	152 8%	61 3%	413 20%	88 4%	660 33%
NET: Top 3 largest causes of death	505 25%	300 15%	941 47%	1735 86%	773 38%	1142 56%	677 33%
Don't know	7 *	6 *	4 *	1 *	6 *	2 *	9 *
Mean	4.59	5.15	3.81	1.97	4.35	3.48	4.63
Standard deviation	1.56	1.49	1.74	1.56	1.95	1.61	2.22
Standard error	0.03	0.03	0.04	0.03	0.04	0.04	0.05

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 2

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Summary table - NET: Top 3 largest causes of death

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Heart disease	1735 86%	862 87% ^b	873 84%	176 78%	255 73%	273 84% ^{cD}	329 91% ^{CDE}	270 91% ^{CDE}	431 93% ^{CDE}	431 75%	603 88% ^I	701 92% ^{IJ}	473 87% ^N	488 87% ⁿ	339 81%	434 87% ⁿ	235 84%	744 84%
Stroke	1142 56%	568 58%	573 55%	101 45%	174 50%	159 49%	211 59% ^{CdE}	196 66% ^{CDEf}	300 64% ^{CDE}	276 48%	370 54% ⁱ	496 65% ^{IJ}	296 55%	349 62% ^{LNO}	224 54%	273 54%	176 63% ^Q	463 52%
Breast cancer	941 47%	440 45%	501 48%	125 55% ^{dFGH}	158 46%	154 47%	161 45%	129 43%	214 46%	284 49%	315 46%	343 45%	233 43%	242 43%	204 49% ^m	262 52% ^{LM}	124 45%	418 47%
Diabetes	773 38%	371 38%	402 39%	90 40%	136 39%	138 42% ^H	133 37%	116 39%	161 34%	226 39%	271 39%	276 36%	226 42% ^O	216 38%	161 39%	170 34%	111 40%	333 38%
Road traffic accident	677 33%	327 33%	351 34%	99 44% ^{FGH}	141 41% ^{FGH}	122 37% ^{GH}	114 32%	77 26%	123 26%	241 42% ^{JK}	236 34% ^K	200 26%	177 33%	174 31%	148 36%	178 36%	85 30%	334 38% ^p
Liver disease	505 25%	248 25%	257 25%	55 24%	111 32% ^{FgH}	90 28% ^H	79 22%	71 24%	98 21%	167 29% ^K	169 25%	169 22%	144 27%	140 25%	101 24%	120 24%	73 26%	232 26%
Kidney disease	300	148	152	33 15%	67 19% ^{EG}	41 13%	52 14%	34 11%	72 15%	101 18%	93 14%	106 14%	77 14%	83 15%	73 17%	68 14%	32 11%	137 15%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 3

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Summary table - NET: Top 3 largest causes of death

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust				
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humber (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)	
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788	
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776	
Heart disease	1735 86%	151 85%	88 87%	1495 86%	69 81%	193 83%	146 86%	145 80%	132 87%h	172 89%H	228 85%	246 87%H	164 92%a	216 EFg	1457 7%	86%	44 73%	233 87%P	272 84%p	1450 86%P	180 83%	1534 86%
Stroke	1142 56%	106 60%	56 56%	979 56%	53 63%	141 60%	91 54%	101 55%	92 61%	103 53%	149 55%	156 55%	93 52%	153 62%o	939 55%	38 63%	143 53%	177 55%	957 57%	121 56%	1005 57%	
Breast cancer	941 47%	79 44%	49 49%	813 47%	43 50%i	104 45%	81 47%i	96 53%lj	57 38%	83 43%	128 47%i	143 51%l	79 44%	103 41%	799 47%	22 36%	127 47%	146 45%	788 47%	94 43%	832 47%	
Diabetes	773 38%	73 41%B	26 26%	674 39%B	33 39%b	94 41%B	59 35%	62 34%	56 37%b	88 45%Bgh	101 37%b	116 41%B	65 37%b	89 36%	662 39%	22 36%	108 40%	128 40%	641 38%	92 42%	667 38%	
Road traffic accident	677 33%	55 31%	30 29%	593 34%	24 28%	70 30%	57 34%	67 37%	54 36%	66 34%	99 37%	91 32%	64 36%	70 28%	581 34%n	16 27%	81 30%	95 30%	578 34%	70 32%	598 34%	
Liver disease	505 25%	45 25%	36 35% Daij KLM	424 24%	22 26%	63 27%	44 26%	47 26%	36 24%	47 24%	63 23%	63 22%	38 21%	66 27%	424 25%	26 43% QrS	69 26%	94 29% s	402 24%	64 30%	430 24%	
Kidney disease	300 15%	26 14%	18 18%l	256 15%	10 12%	34 14%	32 19%L	29 16%	28 18%L	24 12%	39 15%	29 10%	31 17%l	49 20%O	230 14%	14 23%	43 16%	56 17%	236 14%	30 14%	262 15%	

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 4

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Liver disease

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would cause largest no. of deaths	51 3%	19 2%	32 3%	7 3%	12 3%	8 2%	8 2%	4 1%	13 3%	19 3%	15 2%	17 2%	15 3%	19 3% ⁿ	5 1%	12 2%	7 3%	19 2%
2nd largest	176 9%	74 7%	102 10% ^a	20 9% ^f	47 13% ^{FH}	32 10% ^F	16 5%	28 9% ^F	32 7%	67 12% ^{JK}	49 7%	60 8%	41 7%	41 7%	50 12% ^{LM}	45 9%	15 5%	86 10% ^p
3rd largest	278 14%	154 16% ^B	123 12%	28 12%	53 15%	50 15%	55 15% ^h	39 13%	52 11%	81 14%	105 15% ^k	92 12%	88 16% ^N	80 14%	46 11%	63 13%	51 18%	127 14%
4th largest	398 20%	183 19%	216 21%	29 13%	67 19% ^c	58 18%	82 23% ^C	71 24% ^{Ce}	91 20% ^C	96 17%	140 20%	163 21% ⁱ	96 18%	111 20%	85 20%	106 21%	51 18%	165 19%
5th largest	466 23%	229 23%	237 23%	55 24%	63 18%	77 24%	85 23%	73 25% ^d	112 24% ^d	119 21%	161 24%	186 24%	131 24%	119 21%	96 23%	120 24%	62 22%	200 22%
6th largest	430 21%	227 23%	204 20%	63 28% ^{dEFg}	68 19%	62 19%	62 17%	58 20%	116 25% ^{eF}	131 23% ^j	124 18%	175 23% ^J	120 22%	111 20%	93 22%	106 21%	56 20%	194 22%
7th - would cause the smallest no. of deaths	218 11%	99 10%	119 11%	24 11%	38 11%	38 12% ^g	49 14% ^G	22 7%	47 10%	62 11%	87 13% ^K	69 9%	50 9%	79 14% ^{LnO}	41 10%	47 9%	36 13%	93 10%
NET: Top 3 largest causes of death	505 25%	248 25%	257 25%	55 24%	111 32% ^{FgH}	90 28% ^H	79 22%	71 24%	98 21%	167 29% ^K	169 25%	169 22%	144 27%	140 25%	101 24%	120 24%	73 26%	232 26%
Don't know	7 *	3 *	4 *	- -	1 *	1 *	3 1%	2 1%	1 *	1 *	4 1%	3 *	- -	3 1%	1 *	3 1%	1 *	4 *
Mean	4.59	4.63	4.56	4.72 ^D	4.38	4.55	4.69 ^D	4.51	4.71 ^{Dg}	4.51	4.62	4.63	4.57	4.64	4.59	4.57	4.64	4.58
Standard deviation	1.56	1.52	1.59	1.60	1.67	1.59	1.49	1.44	1.52	1.65	1.54	1.49	1.53	1.62	1.54	1.53	1.56	1.57
Standard error	0.03	0.05	0.05	0.11	0.10	0.09	0.08	0.08	0.07	0.08	0.06	0.05	0.06	0.07	0.08	0.07	0.10	0.05

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 5

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Liver disease

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would cause largest no. of deaths	51 3%	3 2%	5 5%KL	43 2%	6 8%AfijKLM	7 3%	6 4%KI	6 3%k	4 3%	5 3%	2 1%	3 1%	3 2%	10 4%	40 2%	4 7%S	9 3%	13 4% _s	38 2%	6 3%	44 2%
2nd largest	176 9%	14 8%	12 12%G	149 9%	8 9%	22 9%	8 5%	13 7%	16 10%g	20 11%g	23 9%	24 9%	15 9%	24 9%	144 8%	8 13%	29 11%	37 11% _s	134 8%	23 10%	150 8%
3rd largest	278 14%	28 16%	19 19%ij	231 13%	9 10%	35 15%	30 17%ij	28 15%	16 11%	21 11%	38 14%	35 13%	20 11%	33 13%	239 14%	13 22%q	31 12%	44 14%	229 14%	35 16%	235 13%
4th largest	398 20%	35 20%	16 16%	348 20%	16 19%	41 18%	27 16%	39 21%	36 24%g	35 18%	46 17%	71 25%bfGk	36 20%	57 23%	322 19%	9 14%	53 20%	61 19%	336 20%	40 19%	355 20%
5th largest	466 23%	37 21%	17 17%	412 24%	20 24%	62 27% _{bk}	45 27% _{bk}	41 23%	39 26%	42 21%	49 18%	63 22%	50 28% _{bK}	56 22%	394 23%	11 18%	74 28%	84 26%	381 23%	53 24%	406 23%
6th largest	430 21%	44 25% _{iL}	25 24% _{iL}	362 21%	13 16%	46 20%	35 20%	34 19%	23 15%	44 23% _L	83 31% _{EF}	42 15%	42 24% _{iL}	43 17%	372 22%	14 23%	55 21%	67 21%	361 21%	39 18%	383 22%
7th - would cause the smallest no. of deaths	218 11%	17 9%	7 7%	194 11%	12 14% _M	19 8%	18 11%	20 11%	17 11%	26 14% _M	27 10%	42 15% _{bFM}	11 6%	23 9%	181 11%	1 1%	15 6%	16 5%	200 12% _{PQR}	19 9%	197 11%
NET: Top 3 largest causes of death	505 25%	45 25%	36 35% _{Daij} KLM	424 24%	22 26%	63 27%	44 26%	47 26%	36 24%	47 24%	63 23%	63 22%	38 21%	66 27%	424 25%	26 43% _{QrS}	69 26%	94 29% _s	402 24%	64 30%	430 24%
Don't know	7 *	1 *	-	7 *	1 1%	2 1%	1 1%	2 1%	1 1%	-	-	* *	-	3 1% _O	4 *	1 2%	2 1%	2 1%	5 *	1 1%	5 *

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 5

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Liver disease

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Mean	4.59	4.62	4.31	4.61b	4.48	4.49	4.61	4.53	4.51	4.68b	4.77Bf	4.64b	4.61	4.42	4.61n	3.96	4.43p	4.34	4.65PQR	4.41	4.62
Standard deviation	1.56	1.52	1.69	1.55	1.76	1.54	1.55	1.56	1.55	1.63	1.51	1.53	1.43	1.58	1.55	1.64	1.51	1.54	1.55	1.57	1.55
Standard error	0.03	0.11	0.16	0.04	0.18	0.10	0.11	0.11	0.12	0.12	0.11	0.09	0.10	0.10	0.04	0.22	0.09	0.09	0.04	0.11	0.04

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 6

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Kidney disease

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would cause largest no. of deaths	34 2%	13 1%	21 2%	6 3%f	13 4%eFgH	3 1%	3 1%	3 1%	4 1%	20 3%JK	6 1%	8 1%	9 2%m	2 *	13 3%M	9 2%M	4 1%	13 1%
2nd largest	79 4%	44 4%	36 3%	8 4%	18 5%	11 3%	12 3%	11 4%	19 4%	26 5%	23 3%	30 4%	19 4%	25 4%	18 4%	17 3%	11 4%	37 4%
3rd largest	187 9%	91 9%	95 9%	19 8%	36 10%	27 8%	38 11%g	19 7%	48 10%g	54 9%	65 9%	68 9%	48 9%	55 10%	41 10%	42 8%	16 6%	87 10%p
4th largest	317 16%	162 16%	155 15%	38 17%d	35 10%	60 18%D	55 15%d	59 20%Dh	69 15%	73 13%	116 17%i	128 17%i	76 14%	87 15%	64 15%	89 18%	44 16%	148 17%
5th largest	460 23%	238 24%	222 21%	48 21%	84 24%	66 20%	87 24%	73 24%	102 22%	133 23%	153 22%	174 23%	139 26%M	114 20%	88 21%	119 24%	61 22%	224 25%
6th largest	526 26%	256 26%	270 26%	45 20%	84 24%	94 29%C	101 28%c	81 27%c	122 26%	129 22%	194 28%l	203 27%	150 28%	157 28%	97 23%	122 24%	81 29%	214 24%
7th - would cause the smallest no. of deaths	415 21%	180 18%	235 23%A	62 27%eFG	76 22%	63 19%	64 18%	47 16%	102 22%g	138 24%Jk	128 19%	149 20%	100 18%	122 22%	92 22%	102 20%	60 22%	160 18%
NET: Top 3 largest causes of death	300 15%	148 15%	152 15%	33 15%	67 19%EG	41 13%	52 14%	34 11%	72 15%	101 18%	93 14%	106 14%	77 14%	83 15%	73 17%	68 14%	32 11%	137 15%
Don't know	6 *	4 *	2 *	- -	1 *	1 *	1 *	3 1%H	- -	1 *	2 *	3 *	- -	2 *	2 *	3 1%	1 *	4 *
Mean	5.15	5.09	5.20	5.19	5.06	5.18	5.15	5.10	5.18	5.11	5.16	5.15	5.15	5.21	5.07	5.13	5.27q	5.06
Standard deviation	1.49	1.46	1.51	1.59	1.65	1.43	1.40	1.38	1.48	1.63	1.41	1.44	1.45	1.46	1.60	1.47	1.44	1.47
Standard error	0.03	0.05	0.05	0.11	0.10	0.08	0.07	0.08	0.07	0.07	0.05	0.05	0.06	0.06	0.09	0.06	0.09	0.05

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 7

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Kidney disease

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbersides (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would cause largest no. of deaths	34 2%	4 2%	1 1%	29 2%	2 2%	6 3%	4 3%	2 1%	2 1%	2 1%	4 1%	4 2%	2 1%	6 2%	21 1%	1 2%	3 1%	5 2%	28 2%	5 2%	28 2%
2nd largest	79 4%	9 5%	4 4%	66 4%	1 1%	6 3%	8 5%	12 6%efl	10 6%efl	7 4%	11 4%	7 3%	5 3%	13 5%	65 4%	2 4%	12 4%	14 4%	64 4%	11 5%	68 4%
3rd largest	187 9%	13 7%	14 14%ajL	160 9%	8 9%	21 9%	19 11%l	15 8%	17 11%	15 7%	25 9%	18 6%	23 13%aL	30 12%o	144 8%	10 17%S	28 10%	38 12%	144 9%	15 7%	166 9%
4th largest	317 16%	26 15%	19 19%	272 16%	13 15%	38 16%	23 14%	26 14%	21 14%	32 17%	44 17%	40 14%	33 19%	37 15%	272 16%	13 22%	44 16%	54 17%	263 16%	44 20%	268 15%
5th largest	460 23%	46 26%F	26 25%F	388 22%	26 30%FJ	36 16%	37 22%	40 22%	31 21%	38 20%	63 23%f	69 24%F	48 27%F	59 24%	381 22%	7 12%	55 20%	62 19%	396 24%p	62 29%u	391 22%
6th largest	526 26%	38 21%	17 17%	471 27%B	24 28%bkM	76 33%ABKM25% m	42 25% m	48 26%bM	45 29%BKM	56 29%BKM	50 19%	101 36%ABGH7% KM	30 7%	61 25%	448 26%	11 19%	69 26%	80 25%	441 26%	61 28%	456 26%
7th - would cause the smallest no. of deaths	415 21%	40 22%l	21 21%	354 20%	11 14%	48 21%	35 21%	39 21%	27 18%	44 22%l	72 27%EIL	42 15%	35 20%	41 17%	362 21%	15 24%	58 22%	71 22%	342 20%	20 9%	392 22%T
NET: Top 3 largest causes of death	300 15%	26 14%	18 18%l	256 15%	10 12%	34 14%	32 19%L	29 16%	28 18%L	24 12%	39 15%	29 10%	31 17%l	49 20%O	230 14%	14 23%	43 16%	56 17%	236 14%	30 14%	262 15%
Don't know	6 *	3 1%DI	-	3 *	-	1 *	1 1%	-	-	1 *	-	-	1 1%	1 *	4 *	-	1 *	1 *	5 *	-	6 *
Mean	5.15	5.13	4.99	5.16	5.10	5.21	5.06	5.13	5.06	5.26m	5.20	5.25m	4.98	4.94	5.20N	4.92	5.14	5.10	5.16	4.89	5.18T
Standard deviation	1.49	1.54	1.48	1.48	1.35	1.53	1.59	1.54	1.54	1.45	1.53	1.33	1.46	1.56	1.46	1.67	1.51	1.55	1.47	1.38	1.50

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 7

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Kidney disease

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Standard error	0.03	0.12	0.14	0.04	0.14	0.10	0.11	0.11	0.12	0.11	0.11	0.08	0.11	0.10	0.04	0.22	0.09	0.09	0.04	0.10	0.04

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 8

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Breast cancer

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would cause largest no. of deaths	183 9%	71 7%	112 11%A	25 11%	43 12%gh	27 8%	30 8%	21 7%	36 8%	69 12%jK	58 8%	57 7%	47 9%	50 9%	36 9%	50 10%	23 8%	86 10%
2nd largest	356 18%	162 16%	193 19%	61 27%DEFGH	60 7%	57 18%	63 17%	44 15%	71 15%	120 21%K	120 17%	115 15%	88 16%	89 16%	76 18%	103 20%lm	42 15%	162 18%
3rd largest	402 20%	206 21%	196 19%	39 17%	55 16%	69 21%	68 19%	64 21%	107 23%D	95 16%	137 20%	171 22%l	98 18%	103 18%	93 22%	109 22%	60 21%	170 19%
4th largest	371 18%	181 18%	190 18%	33 14%	65 19%	56 17%	70 20%	50 17%	97 21%c	98 17%	126 18%	147 19%	123 23%MNO	94 17%	61 15%	93 19%	41 15%	156 18%
5th largest	291 14%	151 15%	140 14%	31 14%	30 9%	54 17%D	61 17%D	52 17%D	63 14%d	61 11%	115 17%l	115 15%l	75 14%	97 17%O	65 16%o	54 11%	50 18%	119 13%
6th largest	265 13%	131 13%	134 13%	25 11%	58 17%E	34 10%	42 12%	44 15%	63 14%	82 14%	76 11%	107 14%	68 13%	88 16%O	57 14%	52 10%	43 15%	115 13%
7th - would cause the smallest no. of deaths	152 8%	82 8%	70 7%	13 6%	37 11%h	29 9%	25 7%	21 7%	28 6%	50 9%	53 8%	49 6%	41 8%	43 8%	28 7%	39 8%	20 7%	76 9%
NET: Top 3 largest causes of death	941 47%	440 45%	501 48%	125 55%dfGH	158 46%	154 47%	161 45%	129 43%	214 46%	284 49%	315 46%	343 45%	233 43%	242 43%	204 49%lm	262 52%LM	124 45%	418 47%
Don't know	4 *	4 *	1 *	- -	- -	- -	1 *	2 1%	1 *	- -	1 *	3 *	1 *	- -	1 *	2 *	- -	3 *
Mean	3.81	3.91B	3.71	3.49	3.86c	3.82c	3.82C	3.95C	3.82C	3.71	3.82	3.87	3.85O	3.95O	3.79	3.62	3.95	3.80
Standard deviation	1.74	1.72	1.76	1.76	1.91	1.74	1.70	1.69	1.65	1.86	1.71	1.67	1.71	1.76	1.74	1.74	1.74	1.79
Standard error	0.04	0.05	0.05	0.12	0.12	0.09	0.09	0.10	0.07	0.09	0.06	0.06	0.07	0.07	0.09	0.08	0.11	0.06

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 9

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Breast cancer

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbersides (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would cause largest no. of deaths	183 9%	12 7%	9 9%	162 9%	13 16% AfghM	20 9%	13 8%	15 8%	14 9%	21 11%	26 10%	27 9%	13 7%	17 7%	159 9%	7 11%	20 7%	26 8%	154 9%	22 10%	158 9%
2nd largest	356 18%	34 19% j	19 19%	303 17%	17 20% j	42 18%	28 17%	41 22% J	19 12%	23 12%	44 16%	60 21% J	29 16%	29 12%	307 18% N	5 8%	47 18%	52 16%	303 18% p	42 19%	305 17%
3rd largest	402 20%	32 18%	22 21%	348 20%	12 14%	42 18%	40 23%	39 22%	24 16%	39 20%	58 21%	57 20%	38 21%	56 23%	332 20%	10 17%	59 22%	68 21%	332 20%	31 14%	369 21% T
4th largest	371 18%	31 17%	14 14%	326 19%	20 24% GK	51 22% GK	22 13%	28 15%	36 24% bGhK KI	49 25% aBGHI KI	34 3%	49 17%	37 21% gk	49 20%	312 18%	14 23%	44 16%	55 17%	311 18%	45 21%	319 18%
5th largest	291 14%	31 17% el	17 17% e	242 14%	7 8%	37 16% e	21 12%	26 14%	20 13%	27 14%	51 19% EL	32 11%	22 12%	40 16%	239 14%	10 17%	37 14%	47 14%	243 14%	25 11%	265 15%
6th largest	265 13%	19 11%	14 14%	232 13%	10 11%	26 11%	33 19% AFHI	21 11%	22 15%	25 13%	34 13%	36 13%	25 14%	42 17% o	210 12%	7 12%	32 12%	39 12%	223 13%	25 11%	235 13%
7th - would cause the smallest no. of deaths	152 8%	17 9%	7 6%	129 7%	6 7%	15 7%	13 8%	11 6%	16 11% j	10 5%	22 8%	20 7%	15 9%	14 6%	135 8%	8 13% s	28 10% s	36 11% S	114 7%	27 13% U	121 7%
NET: Top 3 largest causes of death	941 47%	79 44%	49 49%	813 47%	43 50% i	104 45%	81 47% i	96 53% j	57 38%	83 43%	128 47% i	143 51% l	79 44%	103 41%	799 47%	22 36%	127 47%	146 45%	788 47%	94 43%	832 47%
Don't know	4 *	2 1% D	-	2 *	-	-	-	1 1%	-	-	-	1 *	1 *	1 *	3 *	-	-	-	4 *	1 1%	3 *
Mean	3.81	3.89	3.80	3.80	3.49	3.78	3.96 eh	3.63	4.06 EHL	3.79	3.86	3.67	3.91 e	4.00 o	3.79	4.14	3.89	3.95	3.78	3.89	3.80
Standard deviation	1.74	1.75	1.74	1.74	1.81	1.68	1.79	1.71	1.78	1.66	1.78	1.76	1.72	1.64	1.75	1.83	1.78	1.79	1.73	1.88	1.72

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 9

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Breast cancer

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Standard error	0.04	0.13	0.17	0.04	0.19	0.11	0.13	0.12	0.14	0.12	0.13	0.11	0.12	0.10	0.04	0.24	0.11	0.10	0.04	0.13	0.04

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 10

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Heart disease

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would cause largest no. of deaths	1185 59%	616 62%B	569 55%	97 43%	157 45%	185 57%CD	229 64%CD	211 71%CDEf	306 66%CDE	254 44%	414 60%l	517 68%lJ	309 57%	326 58%	237 57%	313 62%	181 65%Q	488 55%
2nd largest	379 19%	169 17%	210 20%	45 20%	68 19%	61 19%	75 21%G	42 14%	88 19%	113 20%	135 20%	131 17%	122 23%NO	107 19%	65 16%	85 17%	40 14%	172 19%
3rd largest	171 8%	77 8%	94 9%	34 15% ^d EFGH	30 9%	28 9%	25 7%	16 5%	37 8%	64 11% ^j K	53 8%	54 7%	41 8%	56 10%	38 9%	36 7%	15 5%	84 9% ^p
4th largest	91 4%	35 3%	56 5% ^a	20 9% ^e F ^g H	28 8% ^e F ^g H	10 3%	9 2%	8 3%	15 3%	49 8% ^j K	19 3%	23 3%	25 5%	27 5%	22 5%	17 3%	13 5%	48 5%
5th largest	75 4%	28 3%	47 5% ^a	16 7% ^f G ^h	20 6% ^f H	15 5% ^H	8 2%	8 3%	8 2%	36 6% ^j K	23 3%	17 2%	14 3%	24 4%	18 4%	19 4%	13 5%	28 3%
6th largest	62 3%	30 3%	32 3%	8 3% ^G	23 7% ^f G ^h	13 4% ^{G^h}	7 2%	2 1%	8 2%	31 5% ^j K	20 3% ^K	10 1%	14 3%	11 2%	20 5% ^M	18 4%	10 4%	29 3%
7th - would cause the smallest no. of deaths	61 3%	33 3%	28 3%	7 3% ^H	21 6% ^f H	14 4% ^{f^H}	7 2% ^h	9 3% ^H	2 1%	28 5% ^K	21 3% ^k	12 2%	15 3%	14 2%	18 4%	13 3%	7 3%	37 4%
NET: Top 3 largest causes of death	1735 86%	862 87% ^b	873 84%	176 78%	255 73%	273 84% ^c D	329 91% ^{CDE}	270 91% ^{CDE}	431 93% ^{CDE}	431 75%	603 88% ^l	701 92% ^{lJ}	473 87% ^N	488 87% ⁿ	339 81%	434 87% ⁿ	235 84%	744 84%
Don't know	1 *	1 *	- -	- -	- -	- -	- -	- -	1 *	- -	- -	1 *	1 *	- -	- -	- -	- -	- -
Mean	1.97	1.90	2.05 ^a	2.40 ^e F ^g H	2.55 ^e F ^g H	2.09 ^f G ^h	1.72	1.67	1.64	2.49 ^J K	1.90 ^K	1.65	1.92	1.94	2.16 ^L m ^O	1.91	1.92	2.09
Standard deviation	1.56	1.55	1.56	1.65	1.92	1.71	1.31	1.39	1.15	1.82	1.52	1.25	1.48	1.47	1.76	1.55	1.60	1.65
Standard error	0.03	0.05	0.05	0.11	0.12	0.09	0.07	0.08	0.05	0.08	0.06	0.04	0.06	0.06	0.09	0.07	0.10	0.06

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 11

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents
Heart disease

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would cause largest no. of deaths	1185 59%	114 64%beH	54 53%	1017 58%	43 51%	146 63%eH	100 59%	93 51%	93 61%h	107 55%	151 56%	172 61%h	112 63%eH	161 65%O	981 58%	31 52%	164 61%	191 59%	985 59%	118 54%	1049 59%
2nd largest	379 19%	23 13%	19 19%	337 19%a	15 18%	32 14%	37 22%AF	34 19%	27 18%	51 26%AFIL21%af	56 21%	49 17%	35 20%	39 16%	329 19%	11 19%	44 16%	54 17%	323 19%	31 14%	345 19%
3rd largest	171 8%	14 8%	15 15%DaFgjk	142 8%	10 12%g	15 6%	9 5%	18 10%	13 8%	15 8%	21 8%	25 9%	17 9%	16 6%	147 9%	2 3%	26 10%	27 8%	141 8%	31 14%U	139 8%
4th largest	91 4%	7 4%	3 3%	81 5%	7 8%ijM	14 6%k	9 5%	8 5%	4 2%	5 3%	15 5%	16 6%	4 2%	7 3%	79 5%	1 2%	11 4%	12 4%	78 5%	15 7%u	70 4%
5th largest	75 4%	6 4%	3 3%	66 4%	8 9%abFghkIM	5 2%	6 3%	6 3%	7 5%k	12 6%fM	9 3%	10 3%	2 1%	8 3%	64 4%	7 12%QS	10 4%	17 5%	56 3%	6 3%	67 4%
6th largest	62 3%	11 6%DeFJL	6 6%DeFJL	45 3%	1 1%	4 2%	5 3%	10 6%eFJL	4 2%	2 1%	8 3%	5 2%	7 4%	7 3%	51 3%	1 1%	10 4%	11 3%	50 3%	9 4%	53 3%
7th - would cause the smallest no. of deaths	61 3%	2 1%	1 1%	58 3%	1 2%	17 7%ABeGiJLM	4 2%	12 7%ABgJLM	4 3%	2 1%	10 4%	5 2%	1 1%	11 5%	46 3%	7 12%QRS	3 1%	10 3%	50 3%	7 3%	51 3%
NET: Top 3 largest causes of death	1735 86%	151 85%	88 87%	1495 86%	69 81%	193 83%	146 86%	145 80%	132 87%h	172 89%H	228 85%	246 87%H	164 92%aEFgJ7%HK	216 7%	1457 86%	44 73%	233 87%P	272 84%p	1450 86%P	180 83%	1534 86%
Don't know	1 *	-	-	1 *	-	-	-	-	1 *	-	-	-	-	-	1 *	-	-	-	1 *	-	1 *

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 11

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Heart disease

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Mean	1.97	1.94	2.05m	1.97	2.15M	2.05m	1.91	2.34AGI JLM	1.90	1.87	2.03m	1.85	1.74	1.90	1.97	2.56QrS	1.90	2.02	1.96	2.15	1.94
Standard deviation	1.56	1.59	1.50	1.56	1.51	1.80	1.50	1.89	1.53	1.33	1.60	1.40	1.29	1.64	1.53	2.16	1.45	1.64	1.54	1.65	1.54
Standard error	0.03	0.12	0.14	0.04	0.16	0.11	0.11	0.14	0.12	0.10	0.12	0.08	0.09	0.10	0.04	0.29	0.09	0.09	0.04	0.12	0.04

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 12

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Diabetes

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would cause largest no. of deaths	148 7%	58 6%	91 9%A	25 11%Fgh	33 10%F	22 7%	17 5%	18 6%	32 7%	58 10%JK	40 6%	50 7%	38 7%	42 8%	35 8%	32 6%	13 5%	69 8%
2nd largest	291 14%	155 16%	136 13%	33 14%	44 13%	59 18%H	56 16%	42 14%	57 12%	77 13%	115 17%k	99 13%	86 16%	75 13%	63 15%	66 13%	53 19%	126 14%
3rd largest	334 16%	159 16%	175 17%	32 14%	59 17%	57 17%	59 17%	56 19%	71 15%	90 16%	116 17%	127 17%	102 19%o	99 17%	62 15%	71 14%	46 17%	138 16%
4th largest	270 13%	130 13%	141 14%	27 12%	51 15%	41 13%	47 13%	40 14%	64 14%	77 13%	89 13%	104 14%	59 11%	105 19%LNO	50 12%	57 11%	57 21%Q	108 12%
5th largest	299 15%	149 15%	150 14%	30 13%	55 16%	44 13%	55 15%	38 13%	77 17%	85 15%	99 14%	116 15%	87 16%	75 13%	59 14%	78 16%	36 13%	125 14%
6th largest	263 13%	140 14%	122 12%	22 10%	44 13%	37 11%	55 15%c	38 13%	67 14%	65 11%	92 13%	105 14%	73 13%	72 13%	50 12%	67 13%	27 10%	128 14%p
7th - would cause the smallest no. of deaths	413 20%	193 20%	220 21%	59 26%df	62 18%	66 20%	68 19%	61 21%	96 21%	121 21%	134 20%	158 21%	97 18%	93 17%	95 23%IM	127 25%LM	46 16%	189 21%
NET: Top 3 largest causes of death	773 38%	371 38%	402 39%	90 40%	136 39%	138 42%H	133 37%	116 39%	161 34%	226 39%	271 39%	276 36%	226 42%O	216 38%	161 39%	170 34%	111 40%	333 38%
Don't know	6 *	5 *	1 *	- -	- -	1 *	1 *	3 1%	1 *	- -	2 *	4 1%	- -	2 *	1 *	3 1%	1 *	4 *
Mean	4.35	4.37	4.32	4.34	4.24	4.22	4.41	4.36	4.47	4.28	4.32	4.43	4.25	4.22	4.36	4.59LM	4.14	4.40
Standard deviation	1.95	1.92	1.98	2.11	1.94	1.97	1.89	1.93	1.92	2.01	1.93	1.92	1.93	1.87	2.02	1.98	1.83	1.98
Standard error	0.04	0.06	0.06	0.14	0.12	0.11	0.10	0.11	0.08	0.09	0.07	0.07	0.08	0.08	0.11	0.09	0.12	0.07

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 13

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents
Diabetes

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would cause largest no. of deaths	148 7%	7 4%	4 4%	137 8%a	6 7%	13 6%	9 5%	18 10%Ab	13 9%a	18 9%a	26 10%a	20 7%	14 8%	19 7%	122 7%	7 12%	21 8%	27 8%	120 7%	21 10%	124 7%
2nd largest	291 14%	37 21%DBfHI	9 9%	245 14%	10 12%	33 14%	29 17%bHi	17 9%	15 10%	28 15%	40 15%	43 15%h	29 16%h	32 13%	251 15%	7 12%	40 15%	47 15%	244 14%	36 17%	248 14%
3rd largest	334 16%	29 16%	13 13%	292 17%	17 20%	49 21%bGkM	21 12%	27 15%	27 18%	41 21%bGkMI3%	35 19%g	53 19%g	22 13%	38 15%	289 17%	8 13%	48 18%	54 17%	277 16%	35 16%	295 17%
4th largest	270 13%	21 12%	13 13%	237 14%	7 8%	28 12%	31 18%eIJ	30 16%eJ	16 10%	14 7%	51 19%aEflJ	37 13%j	22 12%	30 12%	227 13%	6 9%	33 12%	39 12%	227 13%	21 10%	245 14%
5th largest	299 15%	21 12%	24 17%DAghKLM	255 15%	11 13%	42 18%L	24 14%	26 14%	32 21%AkLm	29 15%	36 13%	32 11%	22 12%	50 20%O	237 14%	17 27%QRS10%	27 10%	42 13%	254 15%Q	21 9%	274 15%T
6th largest	263 13%	16 9%	17 17%aJ	229 13%	12 14%	30 13%	20 12%	22 12%	23 15%j	15 8%	39 15%j	39 14%j	30 17%AJ	32 13%	219 13%	7 11%	31 11%	36 11%	224 13%	32 15%	228 13%
7th - would cause the smallest no. of deaths	413 20%	45 25%FK	21 21%	347 20%	22 26%fk	37 16%	35 21%	41 22%	26 17%	48 25%FK	42 16%	56 20%	39 22%	47 19%	347 20%	10 17%	69 26%S	77 24%	332 20%	50 23%	357 20%
NET: Top 3 largest causes of death	773 38%	73 41%B	26 26%	674 39%B	33 39%b	94 41%B	59 35%	62 34%	56 37%b	88 45%Bgh	101 37%b	116 41%B	65 37%b	89 36%	662 39%	22 36%	108 40%	128 40%	641 38%	92 42%	667 38%
Don't know	6 *	3 1%D	-	4 *	-	1 *	-	1 1%	-	-	-	1 1%	-	1 *	5 *	-	1 *	1 *	6 *	1 1%	5 *
Mean	4.35	4.37	4.77DFjKL	4.32	4.54	4.26	4.37	4.42	4.39	4.26	4.18	4.28	4.43	4.39	4.33	4.28	4.39	4.36	4.34	4.30	4.36

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
*** small base; ** very small base (under 30) ineligible for sig testing**



Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 13

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Diabetes

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Standard deviation	1.95	2.01	1.76	1.95	2.03	1.84	1.92	1.98	1.89	2.06	1.92	1.97	2.02	1.91	1.95	1.96	2.05	2.03	1.93	2.10	1.93
Standard error	0.04	0.15	0.17	0.05	0.21	0.12	0.14	0.14	0.15	0.15	0.14	0.12	0.15	0.12	0.05	0.26	0.12	0.11	0.05	0.15	0.05

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 14

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Stroke

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would cause largest no. of deaths	125 6%	68 7%	57 6%	13 6%	30 9%e	15 5%	19 5%	17 6%	30 6%	43 8%	34 5%	47 6%	34 6%	43 8%	20 5%	28 6%	17 6%	50 6%
2nd largest	589 29%	309 31%b	280 27%	44 20%	71 20%	81 25%	115 32%CD _e	108 36%CD _E	169 36%CD _E	115 20%	197 29% _l	277 36% _l J	147 27%	194 34%LN _O	110 26%	139 28%	95 34% _Q	223 25%
3rd largest	428 21%	191 19%	236 23%a	44 19%	73 21%	63 19%	77 21%	71 24%	100 22%	117 20%	139 20%	171 22%	115 21%	111 20%	95 23%	106 21%	64 23%	190 21%
4th largest	351 17%	177 18%	175 17%	51 22%G _h	66 19% _G	64 20% _G	60 17%	36 12%	75 16%	117 20% _K	124 18% _k	111 15%	106 20% _M	76 13%	85 20% _M	85 17%	40 14%	170 19%
5th largest	222 11%	97 10%	125 12%	26 11%	59 17%FG _H	42 13% _{fgh}	31 9%	23 8%	41 9%	85 15% _{jk}	73 11%	64 8%	52 10%	70 12%	41 10%	59 12%	34 12%	102 11%
6th largest	219 11%	96 10%	123 12%	41 18%DFG _H	32 9%	42 13% _H	41 11% _H	32 11% _h	32 7%	73 13% _K	83 12% _K	64 8%	50 9%	45 8%	59 14%LM	64 13%IM	16 6%	108 12% _P
7th - would cause the smallest no. of deaths	88 4%	49 5%	39 4%	8 3%	16 5%	19 6%	16 4%	9 3%	19 4%	24 4%	35 5%	29 4%	39 7%MN _o	22 4% _n	6 2%	21 4% _N	11 4%	45 5%
NET: Top 3 largest causes of death	1142 56%	568 58%	573 55%	101 45%	174 50%	159 49%	211 59% _{CdE}	196 66% _{CDEf}	300 64% _{CDE}	276 48%	370 54% _i	496 65% _l J	296 55%	349 62%LN _O	224 54%	273 54%	176 63% _Q	463 52%
Don't know	2 *	1 *	1 *	- -	- -	- -	1 *	1 *	- -	- -	1 *	1 *	- -	2 *	- -	- -	1 *	- -
Mean	3.48	3.41	3.54	3.82FG _H	3.61GH	3.73FG _H	3.44 _h	3.25	3.21	3.69 _K	3.57 _K	3.23	3.55 _M	3.28	3.54 _M	3.56 _M	3.26	3.62 _P
Standard deviation	1.61	1.63	1.59	1.61	1.62	1.65	1.62	1.57	1.54	1.62	1.64	1.55	1.66	1.60	1.52	1.62	1.53	1.62
Standard error	0.04	0.05	0.05	0.11	0.10	0.09	0.08	0.09	0.07	0.07	0.06	0.05	0.07	0.07	0.08	0.07	0.10	0.06

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 15

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Stroke

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would cause largest no. of deaths	125 6%	11 6%	11 11%dklm	103 6%	6 8%	18 8%	10 6%	16 9%	9 6%	11 5%	12 4%	13 5%	8 5%	16 6%	104 6%	6 10%	14 5%	19 6%	104 6%	17 8%	105 6%
2nd largest	589 29%	49 28%	34 33%	506 29%	26 31%	79 34%gJ	42 25%	52 29%	53 35%gJ	47 24%	72 27%	85 30%	49 27%	90 36%O	476 28%	21 35%	82 31%	100 31%	482 29%	57 26%	524 30%
3rd largest	428 21%	46 26%Bf	12 12%	370 21%B	21 25%B	43 18%	39 23%B	33 18%	30 20%	45 23%B	65 24%B	58 20%b	36 20%b	48 19%	359 21%	11 18%	47 17%	57 18%	371 22%	47 22%	375 21%
4th largest	351 17%	29 16%	25 25%dfHL	297 17%	14 16%	37 16%	34 20%	26 14%	28 19%	37 19%	48 18%	42 15%	31 17%	37 15%	302 18%	10 17%	51 19%	61 19%	286 17%	35 16%	310 17%
5th largest	222 11%	18 10%	5 5%	199 11%b	6 8%	21 9%	18 10%	26 14%Bi	11 7%	26 14%Bi	29 11%	45 16%BeFIM	16 9%	16 6%	202 12%N	3 5%	34 13%	36 11%	182 11%	27 12%	190 11%
6th largest	219 11%	19 10%	7 7%	193 11%	9 10%	27 12%	17 10%	23 12%	16 11%	18 9%	29 11%	27 10%	27 15%b	30 12%	181 11%	8 13%	25 9%	33 10%	185 11%	27 12%	188 11%
7th - would cause the smallest no. of deaths	88 4%	6 3%	7 7%f	75 4%	3 3%	6 3%	8 5%	7 4%	4 3%	10 5%	14 5%	12 4%	11 6%f	12 5%	73 4%	1 2%	16 6%	16 5%	71 4%	7 3%	80 5%
NET: Top 3 largest causes of death	1142 56%	106 60%	56 56%	979 56%	53 63%	141 60%	91 54%	101 55%	92 61%	103 53%	149 55%	156 55%	93 52%	153 62%o	939 55%	38 63%	143 53%	177 55%	957 57%	121 56%	1005 57%
Don't know	2 *	-	-	2 *	-	1 *	1 1%	-	-	-	-	-	-	1 *	1 *	-	-	-	2 *	-	2 *
Mean	3.48	3.41	3.30	3.49	3.29	3.30	3.55	3.49	3.30	3.60f	3.57	3.53	3.69beF	3.34	3.50	3.19	3.54	3.49	3.48	3.49	3.47
Standard deviation	1.61	1.55	1.71	1.61	1.57	1.60	1.59	1.68	1.54	1.59	1.59	1.59	1.70	1.66	1.60	1.61	1.64	1.64	1.61	1.63	1.61

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 15

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Stroke

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Standard error	0.04	0.12	0.16	0.04	0.16	0.10	0.11	0.12	0.12	0.12	0.11	0.10	0.12	0.10	0.04	0.22	0.10	0.09	0.04	0.11	0.04

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 16

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Road traffic accident

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would cause largest no. of deaths	298 15%	143 14%	155 15%	53 23%FGH	59 17%GH	65 20%GH	54 15%GH	23 8%	44 9%	112 19%K	119 17%K	68 9%	89 17%O	81 14%	71 17%O	56 11%	34 12%	161 18%p
2nd largest	154 8%	75 8%	80 8%	16 7%	40 12%FH	24 7%	23 6%	22 7%	29 6%	56 10%k	47 7%	51 7%	38 7%	33 6%	36 9%	48 10%M	23 8%	80 9%
3rd largest	225 11%	109 11%	116 11%	31 14%	42 12%	33 10%	37 10%	31 11%	50 11%	73 13%	70 10%	82 11%	50 9%	60 11%	41 10%	74 15%Lmn	27 10%	92 10%
4th largest	220 11%	117 12%	103 10%	30 13%	35 10%	37 11%	34 9%	29 10%	56 12%	65 11%	71 10%	85 11%	57 11%	62 11%	49 12%	53 10%	33 12%	89 10%
5th largest	205 10%	93 9%	113 11%	20 9%	36 10%	27 8%	32 9%	27 9%	63 13%Efg	56 10%	59 9%	90 12%j	44 8%	62 11%	49 12%	50 10%	20 7%	86 10%
6th largest	252 12%	103 10%	149 14%A	23 10%	40 11%	44 14%	50 14%	38 13%	57 12%	63 11%	94 14%	95 12%	65 12%	79 14%N	38 9%	70 14%n	44 16%q	96 11%
7th - would cause the smallest no. of deaths	660 33%	344 35%b	316 30%	55 24%	94 27%	95 29%	126 35%Cd	124 42%CDE	166 36%CDe	149 26%	221 32%l	290 38%IJ	195 36%O	185 33%	132 32%	148 29%	96 34%	276 31%
NET: Top 3 largest causes of death	677 33%	327 33%	351 34%	99 44%FGH	141 41%FGH	122 37%GH	114 32%	77 26%	123 26%	241 42%JK	236 34%K	200 26%	177 33%	174 31%	148 36%	178 36%	85 30%	334 38%p
Don't know	9 *	5 *	5 *	- -	1 *	1 *	4 1%	2 1%	1 *	1 *	5 1%	3 *	3 1%	2 *	1 *	3 1%	1 *	6 1%
Mean	4.63	4.65	4.60	4.04	4.28	4.39	4.74CDe	5.12CDEF	4.93CDE	4.19	4.57I	5.00IJ	4.68	4.72	4.47	4.59	4.79Q	4.42
Standard deviation	2.22	2.23	2.20	2.28	2.25	2.31	2.24	2.07	2.05	2.26	2.28	2.06	2.29	2.18	2.26	2.13	2.18	2.30
Standard error	0.05	0.07	0.07	0.15	0.14	0.12	0.11	0.12	0.09	0.10	0.08	0.07	0.10	0.09	0.12	0.09	0.14	0.08

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 17

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Road traffic accident

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would cause largest no. of deaths	298 15%	27 15%	18 18%eF	253 15%	7 9%	22 10%	27 16%f	32 17%eF	18 12%	30 16%f	49 18%eF	41 15%	26 15%	19 8%	269 16%N	4 7%	38 14%	41 13%	255 15%	28 13%	267 15%
2nd largest	154 8%	11 6%	5 5%	138 8%	8 9%	19 8%	17 10%L	13 7%	12 8%	17 9%	23 8%	13 5%	16 9%l	22 9%	124 7%	6 10%	14 5%	19 6%	133 8%	18 8%	136 8%
3rd largest	225 11%	17 9%	7 7%	201 12%	9 10%	29 12%	13 8%	21 12%	24 16%aBG	19 10%	27 10%	36 13%	22 12%	28 11%	188 11%	6 10%	30 11%	35 11%	190 11%	24 11%	196 11%
4th largest	220 11%	27 15%dlm	11 11%	182 10%	8 10%	23 10%	23 14%i	24 13%i	11 7%	21 11%	30 11%	27 10%	15 8%	31 12%	180 11%	8 13%	33 12%	41 13%	177 11%	16 7%	204 11%
5th largest	205 10%	16 9%	9 9%	180 10%	7 8%	29 12%	18 11%	15 8%	12 8%	20 10%	32 12%	30 11%	18 10%	18 7%	177 10%	6 10%	30 11%	35 11%	168 10%	23 10%	178 10%
6th largest	252 12%	29 16%fgk	15 14%	209 12%	17 20%FGKL	24 10%	16 10%	24 13%	19 12%	33 17%fgkl	26 10%	31 11%	18 10%	31 12%	212 13%	13 22%S	46 17%S	57 18%S	194 12%	23 11%	226 13%
7th - would cause the smallest no. of deaths	660 33%	48 27%	37 37%	574 33%	29 34%	87 38%AhJ	54 32%	51 28%	55 36%a	54 28%	82 30%	99 35%a	63 36%	95 38%o	542 32%	17 29%	78 29%	95 29%	557 33%	85 39%u	562 32%
NET: Top 3 largest causes of death	677 33%	55 31%	30 29%	593 34%	24 28%	70 30%	57 34%	67 37%	54 36%	66 34%	99 37%	91 32%	64 36%	70 28%	581 34%n	16 27%	81 30%	95 30%	578 34%	70 32%	598 34%
Don't know	9 *	3 2%Dfjkm	- -	6 *	- -	- -	1 1%	1 1%	1 1%	- -	- -	3 1%	- -	4 2%O	5 *	- -	- -	- -	9 1%	1 1%	8 *
Mean	4.63	4.56	4.79	4.62	4.94hk	4.88HK	4.51	4.41	4.75	4.53	4.40	4.73	4.62	4.95O	4.58	4.90	4.69	4.73	4.60	4.83	4.59
Standard deviation	2.22	2.16	2.30	2.22	2.08	2.11	2.25	2.24	2.21	2.21	2.28	2.21	2.27	2.09	2.23	1.97	2.13	2.10	2.24	2.24	2.21

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 17

Q1. Please rank the following according to how many deaths you think each causes in England and Wales annually, among those of working age (aged 16 to 64). Please rank in order where 1 would cause the largest number of deaths and 7 would cause the smallest number of deaths. If you are unsure, please provide your best guess.

Base: All respondents

Road traffic accident

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Standard error	0.05	0.16	0.22	0.05	0.21	0.13	0.16	0.16	0.18	0.16	0.16	0.13	0.16	0.13	0.05	0.26	0.13	0.12	0.05	0.16	0.05

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 18

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Summary table

	<u>Kidney</u>	<u>Gallbladder</u>	<u>Heart</u>	<u>Lungs</u>	<u>Liver</u>	<u>Skin</u>
Unweighted base	2024	2024	2024	2024	2024	2024
Weighted base	2024	2024	2024	2024	2024	2024
1st - would be of greatest concern	41 2%	56 3%	1461 72%	272 13%	99 5%	70 3%
2nd greatest concern	178 9%	53 3%	314 16%	1069 53%	315 16%	71 3%
3rd greatest concern	568 28%	60 3%	74 4%	341 17%	825 41%	130 6%
4th greatest concern	857 42%	155 8%	56 3%	204 10%	550 27%	173 9%
5th greatest concern	256 13%	844 42%	33 2%	87 4%	170 8%	606 30%
6th would be of least concern	95 5%	826 41%	60 3%	25 1%	38 2%	945 47%
NET: Top 3 greatest concerns	787 39%	169 8%	1849 91%	1681 83%	1239 61%	270 13%
Don't know	29 1%	31 2%	26 1%	27 1%	27 1%	29 1%
Mean	3.70	5.09	1.53	2.42	3.25	5.01
Standard deviation	1.03	1.14	1.14	1.06	1.04	1.30
Standard error	0.02	0.03	0.03	0.02	0.02	0.03

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 19

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

NET: Top 3 greatest concerns

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Heart	1849 91%	907 92%	942 91%	206 91% ^d	295 85%	299 92% ^D	341 95% ^{Dh}	283 95% ^{Dh}	424 91% ^D	502 87%	640 93% ^l	707 93% ^l	490 90%	528 94% ^{lN}	371 89%	460 92%	260 93%	804 91%
Lungs	1681 83%	814 82%	867 84%	189 83%	279 80%	256 78%	315 87% ^{DE}	252 85% ^e	391 84% ^e	467 81%	570 83%	643 84%	457 84% ^N	470 83% ⁿ	327 78%	428 85% ^N	235 84%	723 82%
Liver	1239 61%	613 62%	626 60%	121 53%	192 55%	201 62% ^c	225 63% ^{cd}	200 67% ^{CD}	300 64% ^{CD}	313 54%	426 62% ^l	500 65% ^l	334 62%	345 61%	249 60%	311 62%	167 60%	532 60%
Kidney	787 39%	383 39%	405 39%	85 38%	133 38%	124 38%	143 40%	110 37%	192 41%	218 38%	267 39%	302 40%	217 40%	217 38%	166 40%	188 37%	105 38%	358 40%
Skin	270 13%	136 14%	134 13%	43 19% ^{FGH}	78 22% ^{FGH}	54 17% ^{FGH}	30 8%	21 7%	43 9%	122 21% ^{JK}	84 12% ^K	65 8%	64 12%	84 15%	56 14%	66 13%	40 14%	124 14%
Gallbladder	169 8%	83 8%	86 8%	27 12% ^{FGH}	48 14% ^{FGH}	34 11% ^{FgH}	17 5%	19 6%	23 5%	76 13% ^{JK}	51 7%	42 5%	37 7%	40 7%	54 13% ^{LMO}	38 8%	22 8%	88 10%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 20

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

NET: Top 3 greatest concerns

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humber (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Heart	1849 91%	162 91%	90 89%	1597 92%	73 86%	213 91%	159 94%e	164 90%	138 91%	179 92%	238 88%	271 96%ABEFHjK	162 31%	229 92%	1551 91%	54 90%	243 91%	291 90%	1543 92%	196 90%	1626 92%
Lungs	1681 83%	143 80%	79 78%	1460 84%	75 88%bkm	193 83%	145 85%m	156 86%m	127 84%	164 84%	212 79%	249 88%ABfK	139 78%	191 77%	1430 84%N	47 78%	233 87%	275 85%	1393 83%	169 78%	1488 84%t
Liver	1239 61%	110 62%	69 68%HK	1060 61%	52 61%	145 62%	107 63%	100 55%	93 61%	127 66%hk	149 55%	174 62%	113 63%	167 67%o	1028 61%	36 59%	180 67%s	210 65%	1018 60%	142 66%	1078 61%
Kidney	787 39%	63 35%	44 43%	681 39%	30 36%	89 38%	60 35%	81 44%agjl	65 43%	66 34%	120 44%jl	101 36%	69 39%	99 40%	655 39%	18 29%	93 35%	109 34%	673 40%r	98 45%u	677 38%
Skin	270 13%	24 13%	11 11%	235 13%	15 18%	27 12%	21 12%	19 10%	19 12%	24 12%	51 19%fHL	31 11%	28 16%	33 13%	222 13%	15 25%QRS	27 10%	42 13%	225 13%	18 8%	250 14%T
Gallbladder	169 8%	25 14%DgiJL	9 9%	135 8%	6 7%	20 8%	13 8%	19 10%j	11 7%	9 5%	29 11%J	18 6%	11 6%	20 8%	138 8%	11 19%qS	24 9%	35 11%S	128 8%	27 12%u	137 8%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 21

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Kidney

	Gender		Age									Social Grade				Employment Sector		
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would be of greatest concern	41 2%	23 2%	18 2%	4 2%	4 1%	4 1%	12 3%	4 1%	12 3%	8 1%	16 2%	17 2%	16 3%O	12 2%	8 2%	5 1%	6 2%	21 2%
2nd greatest concern	178 9%	85 9%	93 9%	20 9%	25 7%	36 11%G	30 8%	16 5%	52 11%G	45 8%	66 10%	68 9%	52 10%	48 9%	28 7%	50 10%	20 7%	83 9%
3rd greatest concern	568 28%	275 28%	294 28%	62 27%	103 30%	84 26%	101 28%	90 30%	128 27%	165 29%	185 27%	218 29%	148 27%	157 28%	130 31%	133 27%	79 28%	254 29%
4th greatest concern	857 42%	423 43%	434 42%	67 30%	139 40%C	140 43%C	157 44%C	147 49%CD	207 44%C	207 36%	296 43%I	354 46%I	216 40%	246 44%	164 39%	231 46%In	113 40%	362 41%
5th greatest concern	256 13%	122 12%	134 13%	43 19%dfGH	41 12%	46 14%	46 13%	29 10%	50 11%	84 15%K	92 13%k	79 10%	85 16%MO	62 11%	54 13%	55 11%	44 16%	103 12%
6th would be of least concern	95 5%	49 5%	46 4%	28 12%EFGH	29 8%eFGH	13 4%H	10 3%	8 3%	7 2%	57 10%JK	23 3%	15 2%	15 3%	35 6%L	24 6%L	22 4%	14 5%	53 6%
NET: Top 3 greatest concerns	787 39%	383 39%	405 39%	85 38%	133 38%	124 38%	143 40%	110 37%	192 41%	218 38%	267 39%	302 40%	217 40%	217 38%	166 40%	188 37%	105 38%	358 40%
Don't know	29 1%	11 1%	17 2%	3 1%	6 2%	3 1%	4 1%	3 1%	9 2%	9 2%	7 1%	13 2%	9 2%	5 1%	8 2%	6 1%	4 1%	12 1%
Mean	3.70	3.70	3.70	3.94EFGH	3.80fH	3.70h	3.63	3.69h	3.55	3.86JK	3.67	3.61	3.65	3.72	3.73	3.70	3.76	3.69
Standard deviation	1.03	1.04	1.02	1.22	1.06	1.02	1.02	0.88	0.98	1.13	1.02	0.94	1.05	1.05	1.03	0.98	1.04	1.07
Standard error	0.02	0.03	0.03	0.08	0.07	0.06	0.05	0.05	0.04	0.05	0.04	0.03	0.04	0.04	0.06	0.04	0.07	0.04

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 22

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Kidney

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
		Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would be of greatest concern	41 2%	7 4%GL	1 1%	33 2%	2 2%g	5 2%g	- -	7 4%GL	2 2%	7 4%GL	4 1%	2 1%	4 2%G	7 3%	34 2%	- -	5 2%	5 2%	36 2%	6 3%	34 2%
2nd greatest concern	178 9%	6 3%	14 14%AhjL	158 9%A	6 7%	26 11%Al	15 9%a	12 7%	16 10%A	14 7%	38 14%AHjL	17 6%	15 8%a	26 10%	143 8%	1 2%	22 8%	23 7%	154 9%	30 14%U	147 8%
3rd greatest concern	568 28%	50 28%	29 28%	490 28%	23 27%	58 25%	46 27%	62 34%fJ	47 31%	46 24%	78 29%	81 29%	49 28%	66 27%	478 28%	16 27%	66 24%	81 25%	484 29%	62 28%	495 28%
4th greatest concern	857 42%	77 43%K	47 47%K	733 42%	37 43%k	103 44%K	77 45%K	79 43%K	58 38%	92 47%K	86 32%	131 47%K	70 40%	106 43%	718 42%	28 47%	125 47%	149 46%	701 42%	92 42%	751 42%
5th greatest concern	256 13%	29 16%fH	10 9%	217 12%	11 13%	24 10%	22 13%h	13 7%	23 15%H	21 11%	33 12%	45 16%fH	25 14%h	31 12%	216 13%	11 18%	41 15%	51 16% ^s	201 12%	17 8%	236 13%T
6th would be of least concern	95 5%	7 4%b	- -	88 5%B	6 7%BL	9 4%b	9 5%BL	6 4%b	6 4%b	10 5%BL	28 10%ABFHI	4 2%	10 5%BL	10 4%	84 5%	3 5%	7 3%	10 3%	83 5%	10 4%	86 5%
NET: Top 3 greatest concerns	787 39%	63 35%	44 43%	681 39%	30 36%	89 38%	60 35%	81 44%agjl	65 43%	66 34%	120 44%jl	101 36%	69 39%	99 40%	655 39%	18 29%	93 35%	109 34%	673 40%r	98 45%u	677 38%
Don't know	29 1%	3 2%l	1 1%	25 1%	1 1%	7 3%L	2 1%	3 1%l	1 1%	4 2%L	3 1%	- -	5 3%L	3 1%	24 1%	1 2%	3 1%	3 1%	26 2%	1 *	27 2%
Mean	3.70	3.78Bh	3.51	3.70b	3.80Bh	3.63	3.80BH	3.55	3.66	3.72	3.71	3.76BH	3.72	3.64	3.71	3.96s	3.73	3.78	3.68	3.52	3.72T
Standard deviation	1.03	1.02	0.88	1.03	1.05	1.03	0.96	0.99	1.03	1.06	1.21	0.88	1.07	1.05	1.03	0.86	0.97	0.96	1.04	1.06	1.02
Standard error	0.02	0.08	0.08	0.03	0.11	0.07	0.07	0.07	0.08	0.08	0.09	0.05	0.08	0.06	0.03	0.12	0.06	0.05	0.03	0.07	0.02

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/ef/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 23

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Gallbladder

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would be of greatest concern	56 3%	25 3%	31 3%	8 3% ^{fH}	25 7% ^{eFGH}	11 3% ^{FH}	3 1%	6 2% ^h	3 1%	32 6% ^{JK}	14 2%	9 1%	9 2%	17 3% ^o	23 5% ^{LO}	7 1%	5 2%	34 4%
2nd greatest concern	53 3%	20 2%	32 3%	10 4% ^G	15 4% ^G	10 3% ^G	7 2%	1 *	10 2% ^g	25 4% ^K	17 2%	11 1%	14 3%	8 1%	14 3%	16 3% ^m	7 3%	24 3%
3rd greatest concern	60 3%	38 4% ^b	23 2%	10 4% ^f	9 3%	14 4% ^f	6 2%	11 4%	10 2%	19 3%	20 3%	21 3%	13 2%	15 3%	18 4%	15 3%	10 3%	30 3%
4th greatest concern	155 8%	78 8%	77 7%	22 10% ^g	25 7%	24 7%	28 8%	15 5%	40 9% ^g	46 8%	52 8%	56 7%	39 7%	43 8%	35 8%	38 8%	15 5%	79 9%
5th greatest concern	844 42%	410 42%	434 42%	70 31%	139 40% ^c	120 37%	154 43% ^C	154 52% ^{CDEFh}	207 44% ^{CE}	209 36%	274 40%	361 47% ^{IJ}	207 38%	226 40%	163 39%	247 49% ^{LMN}	107 39%	365 41%
6th would be of least concern	826 41%	406 41%	420 41%	105 46% ^{dG}	130 37%	144 44% ^G	156 43% ^G	105 35%	186 40%	235 41%	300 44% ^K	291 38%	250 46% ^{NO}	251 44% ^{nO}	155 37%	170 34%	131 47% ^Q	341 38%
NET: Top 3 greatest concerns	169 8%	83 8%	86 8%	27 12% ^{FGH}	48 14% ^{FGH}	34 11% ^{FgH}	17 5%	19 6%	23 5%	76 13% ^{JK}	51 7%	42 5%	37 7%	40 7%	54 13% ^{LMO}	38 8%	22 8%	88 10%
Don't know	31 2%	11 1%	20 2%	3 1%	6 2%	4 1%	4 1%	4 1%	10 2%	9 2%	8 1%	14 2%	9 2%	4 1%	10 2% ^m	8 2%	4 1%	13 2%
Mean	5.09	5.09	5.08	5.01	4.84	5.06 ^d	5.23 ^{CDe}	5.13 ^D	5.19 ^{cD}	4.91	5.15 ⁱ	5.16 ⁱ	5.20 ^{NO}	5.15 ^N	4.89	5.05 ⁿ	5.20 ^Q	4.99
Standard deviation	1.14	1.11	1.16	1.30	1.44	1.23	0.93	0.97	0.92	1.39	1.09	0.94	1.06	1.11	1.35	1.03	1.08	1.22
Standard error	0.03	0.04	0.04	0.09	0.09	0.07	0.05	0.06	0.04	0.06	0.04	0.03	0.04	0.05	0.07	0.05	0.07	0.04

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 24

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents
Gallbladder

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would be of greatest concern	56 3%	2 1%	5 5%AjL	49 3%	2 2%	10 4%ajL	4 2%	4 2%	4 3%	2 1%	19 7%AgHJLM	2 1%	3 2%	5 2%	51 3%	4 6%	6 2%	9 3%	46 3%	11 5%u	45 3%
2nd greatest concern	53 3%	14 8%DBFGIJKLm	1 1%	37 2%	3 3%K	5 2%	4 2%	7 4%K	3 2%	3 2%	1 *	6 2%	6 3%K	4 1%	45 3%	3 5%	9 3%	12 4%	38 2%	9 4%	41 2%
3rd greatest concern	60 3%	9 5%jM	3 3%	49 3%	2 2%	5 2%	5 3%	9 5% <i>m</i>	3 2%	3 2%	10 4%	10 4%	2 1%	12 5% <i>o</i>	42 2%	5 8% <i>S</i>	9 3%	14 4%	45 3%	7 3%	51 3%
4th greatest concern	155 8%	7 4%	5 4%	143 8% <i>a</i>	4 4%	15 7%	12 7%	14 7%	12 8%	21 11% <i>Abel</i>	36 13% <i>ABEFg</i>	16 6%	13 8%	12 5%	137 8%	4 7%	17 6%	20 6%	130 8%	17 8%	133 8%
5th greatest concern	844 42%	77 43%	42 41%	725 42%	36 42%	109 47% <i>gk</i>	65 38%	77 42%	73 48% <i>gjK</i>	74 38%	98 36%	118 42%	76 42%	111 44%	700 41%	28 46%	118 44%	143 44%	695 41%	93 43%	737 42%
6th would be of least concern	826 41%	67 38%	45 45%	714 41%	38 45%	82 35%	76 45% <i>f</i>	69 38%	55 36%	86 44% <i>f</i>	104 39%	130 46% <i>aFi</i>	74 42%	102 41%	696 41%	17 28%	107 40%	122 38%	702 42% <i>p</i>	77 36%	741 42%
NET: Top 3 greatest concerns	169 8%	25 14% <i>DgiJL</i>	9 9%	135 8%	6 7%	20 8%	13 8%	19 10% <i>j</i>	11 7%	9 5%	29 11% <i>J</i>	18 6%	11 6%	20 8%	138 8%	11 19% <i>qS</i>	24 9%	35 11% <i>s</i>	128 8%	27 12% <i>u</i>	137 8%
Don't know	31 2%	3 2% <i>l</i>	1 1%	28 2%	1 1%	6 3% <i>L</i>	4 2% <i>L</i>	4 2% <i>L</i>	1 1%	4 2% <i>L</i>	3 1%	-	5 3% <i>L</i>	3 1%	27 2%	-	3 1%	3 1%	29 2%	3 1%	28 2%
Mean	5.09	4.96	5.12	5.10	5.18	5.01	5.16 <i>k</i>	5.02	5.06	5.20 <i>aK</i>	4.90	5.24 <i>AFH</i>	5.16 <i>k</i>	5.14	5.08	4.67	5.08 <i>P</i>	5.00	5.11 <i>P</i>	4.89	5.12 <i>T</i>
Standard deviation	1.14	1.23	1.23	1.12	1.10	1.19	1.10	1.15	1.08	0.98	1.34	0.96	1.05	1.06	1.15	1.41	1.12	1.19	1.12	1.32	1.11

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 24

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Gallbladder

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Standard error	0.03	0.09	0.12	0.03	0.11	0.08	0.08	0.08	0.09	0.07	0.10	0.06	0.08	0.07	0.03	0.19	0.07	0.07	0.03	0.09	0.03

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 25

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Heart

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would be of greatest concern	1461 72%	716 73%	745 72%	172 76%	248 71%	231 71%	274 76% ^H	212 71%	324 69%	421 73%	505 74%	536 70%	406 75% ⁿ	416 74%	287 69%	353 70%	202 72%	650 73%
2nd greatest concern	314 16%	159 16%	154 15%	28 12%	38 11%	48 15%	62 17% ^D	60 20% ^{CDe}	78 17% ^d	67 12%	110 16% ⁱ	137 18% ^l	66 12%	86 15%	76 18% ^L	86 17% ^L	49 18%	124 14%
3rd greatest concern	74 4%	31 3%	43 4%	6 3%	8 2%	20 6% ^{cDF}	5 1%	11 4% ^f	23 5% ^F	14 3%	25 4%	34 4%	18 3%	26 5% ⁿ	8 2%	21 4% ⁿ	8 3%	30 3%
4th greatest concern	56 3%	34 3%	23 2%	3 1%	10 3%	10 3%	12 3%	6 2%	14 3%	13 2%	23 3%	20 3%	17 3%	16 3%	6 1%	17 3% ⁿ	9 3%	26 3%
5th greatest concern	33 2%	15 2%	18 2%	4 2% ^F	12 4% ^{EFg}	3 1%	- -	3 1% ^f	11 2% ^F	16 3% ^J	3 *	14 2% ^J	9 2%	7 1%	12 3%	5 1%	2 1%	13 1%
6th would be of least concern	60 3%	23 2%	37 4%	11 5% ^{FGH}	24 7% ^{eFGH}	11 3% ^F	3 1%	4 1%	8 2%	35 6% ^{JK}	14 2%	11 1%	17 3%	10 2%	19 5% ^M	14 3%	6 2%	33 4%
NET: Top 3 greatest concerns	1849 91%	907 92%	942 91%	206 91% ^d	295 85%	299 92% ^D	341 95% ^{Dh}	283 95% ^{Dh}	424 91% ^D	502 87%	640 93% ^l	707 93% ^l	490 90%	528 94% ^{lN}	371 89%	460 92%	260 93%	804 91%
Don't know	26 1%	9 1%	17 2%	3 1%	6 2%	3 1%	3 1%	2 1%	8 2%	9 2%	6 1%	11 1%	9 2% ^m	3 *	8 2% ^M	5 1%	3 1%	11 1%
Mean	1.53	1.51	1.55	1.53 ^f	1.74 ^{FGh}	1.57 ^F	1.36	1.44	1.55 ^F	1.66 ^{JK}	1.46	1.50	1.51	1.47	1.62 ^m	1.55	1.47	1.55
Standard deviation	1.14	1.09	1.19	1.25	1.50	1.16	0.81	0.90	1.08	1.41	1.00	1.02	1.16	1.01	1.29	1.11	1.01	1.20
Standard error	0.03	0.03	0.04	0.09	0.09	0.06	0.04	0.05	0.05	0.07	0.04	0.04	0.05	0.04	0.07	0.05	0.06	0.04

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 26

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Heart

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would be of greatest concern	1461 72%	130 73%	70 69%	1261 72%	61 72%	171 73%	132 78%JM	128 70%	116 76%jm	130 67%	190 71%	216 77%JM	117 66%	165 66%	1248 74%N	36 60%	169 63%	201 62%	1249 74%PQR	142 65%	1299 73%T
2nd greatest concern	314 16%	26 15%	13 13%	274 16%	10 12%	37 16%	21 12%	31 17%	16 11%	37 19%l	34 13%	50 18%i	37 21%eGIK21%O	51 15%	248 15%	13 21%	61 23%S	72 22%S	240 14%	39 18%	270 15%
3rd greatest concern	74 4%	5 3%	7 7%fhL	62 4%	2 3%	5 2%	6 4%	4 2%	6 4%	12 6%fL	14 5%l	5 2%	7 4%	13 5%	55 3%	6 9%S	13 5%	18 6%S	54 3%	15 7%U	56 3%
4th greatest concern	56 3%	2 1%	3 2%	52 3%	9 11%ABFGH% iJKLM	6 7%GHI	1 1%	5 3%	6 4%GL	4 2%	11 4%gl	3 1%	7 4%gl	7 3%	46 3%	1 2%	6 2%	7 2%	48 3%	8 4%	48 3%
5th greatest concern	33 2%	3 2%	1 1%	29 2%	- -	6 3%	3 2%	6 3%l	2 1%	2 1%	5 2%	2 1%	3 2%	7 3%	25 1%	1 2%	14 5%S	16 5%S	17 1%	3 1%	31 2%
6th would be of least concern	60 3%	9 5%flm	7 7%DFLM	45 3%	2 2%	4 2%	6 3%	5 3%	5 3%	5 2%	13 5%lm	5 2%	2 1%	3 1%	53 3%	4 6%Q	3 1%	6 2%	52 3%q	10 5%	48 3%
NET: Top 3 greatest concerns	1849 91%	162 91%	90 89%	1597 92%	73 86%	213 91%	159 94%e	164 90%	138 91%	179 92%	238 88%	271 96%ABEF HijKM	162 91%	229 92%	1551 91%	54 90%	243 91%	291 90%	1543 92%	196 90%	1626 92%
Don't know	26 1%	3 2%l	1 1%	22 1%	1 1%	4 2%L	2 1%	3 1%l	1 1%	4 2%L	3 1%	- -	5 3%L	2 1%	22 1%	- -	2 1%	2 1%	24 1%	1 *	24 1%
Mean	1.53	1.56l	1.72dgL	1.52	1.60l	1.48	1.44	1.58L	1.52	1.55l	1.66L	1.36	1.55l	1.57	1.51	1.82s	1.67S	1.70S	1.49	1.71U	1.51
Standard deviation	1.14	1.25	1.40	1.11	1.17	1.06	1.10	1.19	1.17	1.06	1.32	0.88	1.01	1.06	1.14	1.38	1.15	1.21	1.11	1.28	1.11
Standard error	0.03	0.09	0.13	0.03	0.12	0.07	0.08	0.09	0.09	0.08	0.10	0.05	0.07	0.07	0.03	0.18	0.07	0.07	0.03	0.09	0.03

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/ef/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 27

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Lungs

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would be of greatest concern	272 13%	144 15%	128 12%	28 13%	28 8%	40 12%	45 13% ^d	51 17% ^D	80 17% ^{Def}	56 10%	85 12%	131 17% ^{IJ}	64 12%	63 11%	56 13%	90 18% ^{LM}	37 13%	104 12%
2nd greatest concern	1069 53%	516 52%	552 53%	132 58% ^H	196 56% ^H	164 50%	207 58% ^{eH}	157 53% ^h	212 45%	328 57% ^K	371 54% ^K	369 48%	301 56% ⁿ	310 55% ⁿ	201 48%	256 51%	158 57%	467 53%
3rd greatest concern	341 17%	153 16%	187 18%	28 12%	55 16%	52 16%	62 17%	44 15%	99 21% ^{CdeG}	83 14%	114 17%	144 19% ⁱ	92 17%	97 17%	70 17%	82 16%	41 15%	153 17%
4th greatest concern	204 10%	108 11%	96 9%	22 10%	37 11%	36 11%	28 8%	34 12%	47 10%	59 10%	64 9%	81 11%	52 10%	51 9%	52 12%	50 10%	30 11%	94 11%
5th greatest concern	87 4%	43 4%	44 4%	11 5%	17 5%	24 7% ^{FGH}	10 3%	8 3%	16 3%	28 5%	34 5% ^k	24 3%	19 4%	30 5% ^o	24 6% ^O	14 3%	6 2%	45 5% ^p
6th would be of least concern	25 1%	12 1%	12 1%	2 1%	9 3% ^{fg}	7 2% ^g	2 1%	1 *	4 1%	11 2% ^k	9 1%	5 1%	5 1%	10 2%	4 1%	6 1%	5 2%	12 1%
NET: Top 3 greatest concerns	1681 83%	814 82%	867 84%	189 83%	279 80%	256 78%	315 87% ^{DE}	252 85% ^e	391 84% ^e	467 81%	570 83%	643 84%	457 84% ^N	470 83% ⁿ	327 78%	428 85% ^N	235 84%	723 82%
Don't know	27 1%	11 1%	17 2%	3 1%	6 2%	3 1%	5 1%	2 1%	8 2%	9 2%	8 1%	11 1%	9 2%	3 1%	10 2% ^M	5 1%	3 1%	13 1%
Mean	2.42	2.41	2.42	2.38	2.55 ^{FGH}	2.57 ^{cFGH}	2.31	2.30	2.39	2.48 ^k	2.44	2.35	2.39	2.47 ^O	2.51 ^O	2.31	2.37	2.48
Standard deviation	1.06	1.08	1.04	1.05	1.11	1.19	0.93	0.99	1.05	1.09	1.07	1.03	1.00	1.09	1.12	1.04	1.03	1.09
Standard error	0.02	0.03	0.03	0.07	0.07	0.06	0.05	0.06	0.05	0.05	0.04	0.04	0.04	0.05	0.06	0.05	0.07	0.04

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 28

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Lungs

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West	Yes	No	Yes - me	Yes - my family/friends	Yes - me/family/friends	No	Yes	No
		(a)	(b)	(d)	(e)	(f)	(g)	(h)	(i)	(j)	(k)	(l)	(m)	(n)	(o)	(p)	(q)	(r)	(s)	(t)	(u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would be of greatest concern	272 13%	20 11%	10 10%	241 14%	12 14%	29 13%	21 12%	30 16%	21 14%	28 14%	34 12%	38 14%	29 16%	38 15%	220 13%	12 19%	58 22% ^S	68 21% ^S	203 12%	22 10%	243 14%
2nd greatest concern	1069 53%	106 60% ^{dkm}	53 52%	909 52%	48 56% ^m	120 51%	99 58% ^M	92 51%	77 51%	103 53% ^m	134 50%	160 57% ^M	76 43%	110 44%	926 55% ^N	24 40%	126 47%	149 46%	909 54% ^{ppqR}	101 47%	955 54% ^t
3rd greatest concern	341 17%	17 9%	15 15%	309 18% ^A	15 18% ^a	44 19% ^A	26 15%	34 19% ^A	28 19% ^A	33 17% ^a	44 17% ^a	51 18% ^A	34 19% ^A	43 17%	284 17%	12 19%	49 18%	59 18%	281 17%	46 21%	289 16%
4th greatest concern	204 10%	26 15% ^{DHJK}	13 13%	165 9%	7 8%	24 10%	16 10%	13 7%	21 14% ^{hk}	17 9%	20 7%	23 8%	24 13% ^{hk}	44 18% ^O	152 9%	8 14%	24 9%	32 10%	171 10%	23 11%	178 10%
5th greatest concern	87 4%	5 3%	8 8% ^{dafGH}	73 4% ^{IL}	2 3%	7 3%	4 2%	4 2%	3 2%	8 4%	28 11% ^{AeFGH}	8 3% ^{IJLM}	7 4%	4 2%	75 4% ⁿ	2 3%	6 2%	8 2%	76 5%	20 9% ^U	66 4%
6th would be of least concern	25 1%	1 *	* *	24 1%	- -	5 2% ^l	3 2% ^l	5 2% ^l	- -	1 *	6 2% ^l	1 *	4 2% ^l	6 2% ^o	18 1%	3 4% ^S	3 1%	6 2%	18 1%	3 1%	19 1%
NET: Top 3 greatest concerns	1681 83%	143 80%	79 78%	1460 84%	75 88% ^{bkm}	193 83%	145 85% ^m	156 86% ^m	127 84%	164 84%	212 79%	249 88% ^{ABfK}	139 78% ^M	191 77%	1430 84% ^N	47 78%	233 87%	275 85%	1393 83%	169 78%	1488 84% ^t
Don't know	27 1%	3 2%	1 1%	24 1%	1 1%	4 2% ^l	2 1%	4 2% ^l	1 1%	4 2% ^l	3 1%	1 *	5 3% ^L	3 1%	23 1%	- -	2 1%	2 1%	26 2%	2 1%	25 1%
Mean	2.42	2.39	2.58 ^{eL}	2.41	2.29	2.45	2.37	2.35	2.38	2.35	2.60 ^{egh}	2.31	2.52 ^L	2.53 ^o	2.40	2.56	2.26	2.32	2.43 ^Q	2.66 ^U	2.39
Standard deviation	1.06	1.00	1.12	1.07	0.91	1.07	1.02	1.08	0.96	1.01	1.25	0.92	1.18	1.15	1.04	1.29	1.04	1.10	1.05	1.17	1.04
Standard error	0.02	0.08	0.11	0.03	0.09	0.07	0.07	0.08	0.08	0.07	0.09	0.06	0.09	0.07	0.03	0.17	0.06	0.06	0.03	0.08	0.02

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing



Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 29

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Liver

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would be of greatest concern	99 5%	42 4%	57 5%	3 1%	14 4%	22 7% C	17 5% C	17 6% C	27 6% C	17 3%	39 6% l	44 6% l	26 5%	29 5%	22 5%	23 5%	12 4%	42 5%
2nd greatest concern	315 16%	164 17%	150 15%	21 9%	47 13%	53 16% C	45 12%	54 18% Cf	95 20% CDF	68 12%	98 14%	149 19% lJ	83 15%	86 15%	67 16%	78 16%	34 12%	144 16%
3rd greatest concern	825 41%	407 41%	418 40%	97 43%	131 38%	126 39%	163 45% dH	129 43%	178 38%	228 40%	289 42%	307 40%	225 42%	231 41%	159 38%	210 42%	120 43%	346 39%
4th greatest concern	550 27%	258 26%	292 28%	75 33% egh	102 29%	81 25%	102 28%	75 25%	116 25%	176 31% k	183 27%	191 25%	156 29%	147 26%	116 28%	132 26%	87 31%	238 27%
5th greatest concern	170 8%	83 8%	87 8%	23 10%	32 9%	34 10% g	24 7%	18 6%	39 8%	55 9%	58 9%	57 7%	37 7%	59 10% L	29 7%	46 9%	21 7%	81 9%
6th would be of least concern	38 2%	23 2%	15 1%	6 2% H	17 5% FGH	7 2% H	5 1%	2 1%	2 *	22 4% JK	12 2% K	4 1%	7 1%	11 2%	13 3% l	7 1%	1 *	26 3% P
NET: Top 3 greatest concerns	1239 61%	613 62%	626 60%	121 53%	192 55%	201 62% c	225 63% cd	200 67% CD	300 64% CD	313 54%	426 62% l	500 65% l	334 62%	345 61%	249 60%	311 62%	167 60%	532 60%
Don't know	27 1%	10 1%	17 2%	3 1%	6 2%	3 1%	3 1%	2 1%	9 2%	9 2%	6 1%	12 2%	9 2% m	3 *	8 2% M	6 1%	3 1%	11 1%
Mean	3.25	3.25	3.24	3.49 EFGH	3.41 efGH	3.22	3.25 gh	3.10	3.11	3.44 JK	3.24 K	3.11	3.22	3.27	3.25	3.24	3.27	3.29
Standard deviation	1.04	1.05	1.04	0.93	1.11	1.12	0.97	0.98	1.03	1.05	1.04	1.01	0.99	1.07	1.09	1.02	0.95	1.09
Standard error	0.02	0.03	0.03	0.06	0.07	0.06	0.05	0.06	0.05	0.05	0.04	0.04	0.04	0.04	0.06	0.05	0.06	0.04

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 30

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Liver

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would be of greatest concern	99 5%	5 3%	12 11% DAEFG HiKL	82 5%	2 2%	9 4%	7 4%	5 3%	7 5%	14 7% aeh	10 4%	14 5%	15 8% AefHk	23 9% O	67 4%	3 6%	21 8% S	24 7% S	73 4%	24 11% U	74 4%
2nd greatest concern	315 16%	21 12%	16 16%	277 16%	11 13%	38 16%	26 15%	33 18%	30 20% aj	23 12%	46 17%	42 15%	29 16%	52 21% O	249 15%	17 28% qS	43 16%	57 18%	257 15%	34 16%	271 15%
3rd greatest concern	825 41%	83 47% HiK	41 41%	701 40%	39 46% hk	98 42%	74 44% h	63 34%	56 37%	90 46% HiK	93 35%	118 42%	69 39%	92 37%	712 42%	16 26%	116 43% P	128 40% p	688 41% P	84 39%	733 41%
4th greatest concern	550 27%	49 27%	24 24%	477 27%	23 27%	63 27%	44 26%	53 29%	42 28%	43 22%	82 31% j	85 30% j	42 24%	57 23%	469 28%	15 25%	62 23%	76 23%	471 28%	57 26%	484 27%
5th greatest concern	170 8%	11 6%	6 6%	153 9%	6 7%	18 8%	15 9%	22 12% al	13 8%	16 8%	30 11%	19 7%	13 8%	19 8%	143 8%	6 11%	17 6%	23 7%	144 9%	14 7%	154 9%
6th would be of least concern	38 2%	6 3%	1 1%	31 2%	3 4%	2 1%	2 1%	4 2%	3 2%	4 2%	4 2%	3 1%	5 3%	4 2%	34 2%	3 6% S	8 3%	12 4% S	26 2%	3 1%	34 2%
NET: Top 3 greatest concerns	1239 61%	110 62%	69 68% Hk	1060 61%	52 61%	145 62%	107 63%	100 55%	93 61%	127 66% hk	149 55%	174 62%	113 63%	167 67% o	1028 61%	36 59%	180 67% s	210 65%	1018 60%	142 66%	1078 61%
Don't know	27 1%	3 2% l	1 1%	23 1%	1 1%	5 2% L	2 1%	3 1% l	1 1%	4 2% L	3 1%	-	5 3% L	2 1%	23 1%	-	2 1%	2 1%	25 1%	1 *	25 1%
Mean	3.25	3.32B	3.00	3.25B	3.36B	3.22b	3.25b	3.38Bm	3.22	3.18	3.34B	3.22b	3.15	3.03	3.28N	3.24	3.13	3.16	3.26q	3.06	3.27T
Standard deviation	1.04	0.98	1.10	1.04	1.00	0.98	1.00	1.08	1.08	1.07	1.06	0.98	1.14	1.13	1.02	1.29	1.11	1.15	1.02	1.12	1.03
Standard error	0.02	0.07	0.11	0.03	0.10	0.06	0.07	0.08	0.09	0.08	0.08	0.06	0.08	0.07	0.02	0.17	0.07	0.06	0.03	0.08	0.02

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/ef/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 31

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Skin

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
1st - would be of greatest concern	70 3%	28 3%	42 4%	9 4%fg	23 6%FGH	15 5%FG	6 2%	4 1%	13 3%	32 6%jk	21 3%	17 2%	12 2%	26 5%L	12 3%	20 4%	15 5%	26 3%
2nd greatest concern	71 3%	33 3%	37 4%	13 6%Fgh	21 6%FgH	13 4%f	5 1%	7 2%	12 3%	34 6%JK	18 3%	19 2%	16 3%	22 4%	22 5%O	11 2%	6 2%	34 4%
3rd greatest concern	130 6%	75 8%b	55 5%	21 9%fGH	35 10%FGH	26 8%GH	19 5%	10 3%	19 4%	56 10%jk	45 7%K	29 4%	36 7%	36 6%	22 5%	35 7%	18 7%	64 7%
4th greatest concern	173 9%	76 8%	97 9%	35 16%DeFGH	29 8%	32 10%g	28 8%	16 5%	32 7%	65 11%K	60 9%	49 6%	53 10%O	57 10%O	34 8%	28 6%	22 8%	77 9%
5th greatest concern	606 30%	304 31%	302 29%	73 32%	100 29%	96 29%	121 34%	82 28%	134 29%	173 30%	217 32%	216 28%	176 32%O	177 31%o	126 30%	128 25%	95 34%	268 30%
6th would be of least concern	945 47%	461 47%	484 47%	73 32%	134 38%	141 43%C	176 49%CD	174 59%CDEF	248 53%CDE	206 36%	317 46%I	422 55%IJ	240 44%	242 43%	190 46%	273 54%LMN	118 42%	405 46%
NET: Top 3 greatest concerns	270 13%	136 14%	134 13%	43 19%FGH	78 22%FGH	54 17%FGH	30 8%	21 7%	43 9%	122 21%JK	84 12%K	65 8%	64 12%	84 15%	56 14%	66 13%	40 14%	124 14%
Don't know	29 1%	11 1%	19 2%	3 1%	6 2%	3 1%	6 2%	3 1%	9 2%	9 2%	8 1%	12 2%	9 2%	4 1%	10 2% ^m	7 1%	4 1%	13 1%
Mean	5.01	5.02	5.00	4.64	4.65	4.87	5.20CDE	5.34CDE	5.20CDE	4.65	5.04I	5.25IJ	5.03	4.90	4.99	5.13M	4.93	4.99
Standard deviation	1.30	1.27	1.33	1.38	1.54	1.40	1.07	1.06	1.19	1.48	1.25	1.14	1.20	1.37	1.32	1.31	1.36	1.29
Standard error	0.03	0.04	0.04	0.09	0.10	0.08	0.05	0.06	0.05	0.07	0.05	0.04	0.05	0.06	0.07	0.06	0.09	0.05

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 32

Q2. Please rank the following in terms of how concerned you would be, if at all, if you were told you had a problem with each. Please rank in order where 1 would be of greatest concern and 6 would be of least concern.

Base: All respondents

Skin

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humber (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
1st - would be of greatest concern	70 3%	11 6%Dfl	3 3%	56 3%	6 7%fl	5 2%	5 3%i	6 3%i	1 *	8 4%l	10 4%i	9 3%i	6 3%i	9 3%	55 3%	5 9%qS	8 3%	13 4%	54 3%	12 5%	56 3%
2nd greatest concern	71 3%	2 1%	3 3%	66 4%a	7 8%AFghL	3 1%	5 3%	5 3%	8 5%AF	9 5%AF	14 5%AF	6 2%	10 6%AFI	4 1%	64 4%n	3 5%	5 2%	8 2%	62 4%	3 1%	67 4%
3rd greatest concern	130 6%	11 6%	6 6%	113 6%	2 3%	19 8%j	11 6%	8 4%	10 7%	7 4%	28 10%ehJ	16 6%	12 7%	21 8%	103 6%	6 11%	14 5%	21 6%	109 6%	3 1%	127 7%T
4th greatest concern	173 9%	14 8%	9 9%	150 9%	5 6%	15 7%	17 10%	15 8%	12 8%	13 7%	31 12%	23 8%	17 9%	19 8%	151 9%	4 6%	32 12%S	36 11%S	135 8%	18 8%	155 9%
5th greatest concern	606 30%	50 28%	33 33%	523 30%	29 34%	62 27%	59 35%fi	56 31%	37 25%	68 35%fi	73 27%	89 32%	49 28%	74 30%	515 30%	12 20%	71 26%	80 25%	524 31%R	69 32%	526 30%
6th would be of least concern	945 47%	87 49%	47 46%	812 47%	35 41%	124 53%eGjK	71 42%	89 49%	81 53%eGjK	84 43%	111 41%	137 49%	79 44%	119 48%	784 46%	30 49%	136 51%	163 51%	772 46%	111 51%	818 46%
NET: Top 3 greatest concerns	270 13%	24 13%	11 11%	235 13%	15 18%	27 12%	21 12%	19 10%	19 12%	24 12%	51 19%fHL	31 11%	28 16%	33 13%	222 13%	15 25%QRS	27 10%	42 13%	225 13%	18 8%	250 14%T
Don't know	29 1%	3 2%	1 1%	26 1%	1 1%	5 2%L	2 1%	4 2%i	2 2%	4 2%i	3 1%	1 *	5 3%L	3 1%	25 1%	-	2 1%	2 1%	28 2%	2 1%	27 2%
Mean	5.01	5.00	5.07	5.01	4.77	5.18EKm	4.98	5.11eK	5.15eK	4.99	4.79	5.10eK	4.91	5.06	5.01	4.71	5.11p	5.03	5.01	5.15	4.99
Standard deviation	1.30	1.40	1.22	1.30	1.57	1.19	1.24	1.25	1.19	1.33	1.40	1.22	1.38	1.26	1.30	1.68	1.22	1.33	1.29	1.28	1.30
Standard error	0.03	0.11	0.12	0.03	0.16	0.08	0.09	0.09	0.10	0.10	0.10	0.07	0.10	0.08	0.03	0.22	0.07	0.07	0.03	0.09	0.03

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/ef/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 33

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

First mention

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
NET: Any mentions of Alcohol	1472 73%	706 71%	766 74%	133 59%	246 71%C	224 69%C	275 76%CE	235 79%CDE	359 77%CdE	379 66%	499 73%I	594 78%J	407 75%O	415 73%	306 73%	345 69%	200 72%	631 71%
Alcohol/ drinking alcohol	1014 50%	473 48%	541 52%a	87 38%	177 51%C	149 46%	193 54%Ce	166 56%CE	242 52%C	264 46%	342 50%	408 53%I	269 50%	284 50%	217 52%	243 48%	147 53%	448 51%
Excessive alcohol consumption	327 16%	172 17%	155 15%	28 13%	39 11%	52 16%	54 15%	51 17%d	103 22%CDEF	67 12%	105 15%	154 20%J	94 17%	99 18%	62 15%	72 14%	27 10%	125 14%
Alcoholism/ alcohol addiction/ abuse	131 6%	61 6%	70 7%	18 8%H	30 9%H	23 7%H	28 8%H	18 6%H	14 3%	48 8%K	51 7%K	32 4%	44 8%	31 6%	26 6%	30 6%	27 10%	58 7%
Smoking/ cigarettes	46 2%	21 2%	25 2%	13 6%DFGH	7 2%	9 3%f	3 1%	5 2%	9 2%	20 3%k	12 2%	13 2%	6 1%	16 3%I	13 3%I	11 2%	8 3%	14 2%
A poor diet/ eating bad food (non-specific/ other)	32 2%	20 2%	11 1%	- -	8 2%cg	8 3%Cfg	2 1%	1 *	12 3%CFG	8 1%	10 2%	13 2%	8 2%	8 1%	10 2%	6 1%	5 2%	13 1%
Fatty food	13 1%	5 *	8 1%	1 1%	3 1%	3 1%	3 1%	2 1%	1 *	5 1%	5 1%	3 *	5 1%	2 *	2 1%	3 1%	3 1%	5 1%
Hepatitis	11 1%	4 *	7 1%	* *	- -	2 1%	* *	3 1%	5 1%df	* *	3 1%I	8 1%I	2 *	5 1%	2 *	3 1%	2 1%	2 *
Obesity/ being overweight	10 *	3 *	7 1%	1 *	2 *	2 1%	3 1%	1 *	1 *	2 *	5 1%	2 *	5 1%m	1 *	1 *	3 1%	2 1%	5 1%
Drug use/ abuse	5 *	2 *	4 *	- -	1 *	1 *	1 *	1 *	1 *	1 *	2 *	2 *	2 *	- -	2 1%	1 *	- -	3 *
An infection	5 *	1 *	4 *	3 1%de	- -	- -	* *	1 *	1 *	3 *	* *	2 *	4 1%o	1 *	- -	- -	1 *	3 *
Diabetes	3 *	1 *	3 *	- -	- -	1 *	1 *	1 *	* *	- -	2 *	1 *	1 *	1 *	- -	1 *	2 1%Q	* *

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 33
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
First mention

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Other condition/ illness	3*	1*	2*	* *	- -	- -	1* *	- -	2* *	* *	1* *	2* *	- -	2* *	- -	1* *	- -	1* *
Medication/ overuse of medication	2*	2*	- -	1* *	- -	- -	- -	1* *	1* *	1* *	- -	1* *	- -	1* *	1* *	- -	- -	1* *
Cancer	2*	2*	- -	* *	- -	- -	- -	1* *	1* *	* *	- -	2* *	* *	- -	2* *	- -	- -	2* *
Blood/ blood disorder/ infection	2*	1*	1*	- -	- -	- -	- -	- -	2* *	- -	- -	2* *	- -	- -	- -	2* *	- -	- -
High cholesterol	2*	1*	2*	- -	- -	1* *	1* *	- -	1* *	- -	1* *	1* *	- -	1* *	- -	2* *	- -	1* *
Sugar	1*	- -	1* *	- -	1* *	- -	- -	- -	- -	1* *	- -	- -	1* *	- -	- -	- -	- -	1* *
Genetics/ family history	1*	1*	- -	- -	- -	- -	- -	* *	1* *	- -	- -	1* *	1* *	- -	- -	- -	- -	- -
Other answers	5*	3*	2*	- -	1* *	1* *	3* 1%H	- -	- -	1* *	4* 1%K	- -	1* *	2* *	1* *	1* *	1* *	5* 1%
None/ Don't know	409 20%	213 22%	195 19%	73 32% DEFGH	79 3% GH	74 23% GH	66 18%	45 15%	71 15%	152 26% JK	140 20% K	116 15%	97 18%	110 19%	76 18%	125 25% LMN	54 19%	202 23%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 34

Q3. Can you name three things that you think would make you more likely to develop liver disease?**Base: All respondents****First mention**

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humber (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/ friends (q)	Yes - me/ family/ friends (r)	No (s)	Yes (t)	No (u)
		Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
NET: Any mentions of Alcohol	1472	136	72	1265	53	176	122	131	111	150	179	208	134	188	1237	41	218	254	1207	160	1293
	73%	76%Ek	71%	72%	63%	75%Ek	72%	72%	73%	77%EK	67%	74%e	75%e	76%	73%	68%	81%pS	79%S	72%	74%	73%
Alcohol/ drinking alcohol	1014	89	57	868	35	123	92	86	77	111	117	134	92	126	864	27	151	173	832	108	889
	50%	50%	57%ek	50%	41%	53%ek	54%ek	47%	50%	57%EhKl	44%	48%	52%	51%	51%	45%	56%e	54%	49%	50%	50%
Excessive alcohol consumption	327	31	11	285	17	38	19	31	29	27	38	56	31	49	266	9	49	56	269	32	295
	16%	18%	10%	16%	19%g	16%	11%	17%	19%bg	14%	14%	20%B	17%	20%	16%	14%	18%	17%	16%	15%	17%
Alcoholism/ alcohol addiction/ abuse	131	16	4	111	2	15	11	14	5	12	24	18	11	14	107	6	18	24	106	20	110
	6%	9%ei	4%	6%	2%	6%	6%	8%e	4%	6%	9%ei	6%	6%	6%	6%	10%	7%	7%	6%	9%	6%
Smoking/ cigarettes	46	1	2	43	1	2	3	7	7	2	11	6	3	3	40	2	8	10	35	10	36
	2%	*	2%	2%	1%	1%	2%	4%Afj	5%AFJ	1%	4%af	2%	2%	1%	2%	4%	3%	3%	2%	4%u	2%
A poor diet/ eating bad food (non-specific/ other)	32	1	4	28	-	2	1	6	4	-	3	9	3	8	24	2	6	8	24	2	30
	2%	*	3%aj	2%	-	1%	1%	3%aj	3%j	-	1%	3%afj	2%	3%o	1%	3%	2%	3%	1%	1%	2%
Fatty food	13	1	-	12	-	2	*	1	2	4	-	1	1	2	9	1	4	4	7	1	12
	1%	1%	-	1%	-	1%	*	1%	1%	2%k	-	*	*	1%	1%	1%	1%e	1%e	*	*	1%
Hepatitis	11	1	1	9	1	-	-	-	1	*	2	4	1	2	8	-	3	3	8	1	10
	1%	1%	1%	1%	1%	-	-	-	1%	*	1%	1%	1%	1%	*	-	1%	1%	*	*	1%
Obesity/ being overweight	10	-	-	10	-	4	1	1	1	-	-	2	1	-	10	-	1	1	9	4	4
	*	-	-	1%	-	2%k	1%	*	1%	-	-	1%	*	-	1%	-	*	*	1%	2%u	*
Drug use/ abuse	5	2	-	3	-	-	1	1	-	-	-	1	-	-	5	-	2	2	3	-	5
	*	1%D	-	*	-	-	*	1%	-	-	-	*	-	-	*	-	1%e	1%	*	-	*

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 34
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
First mention

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotlan d (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshi re & Humber s (g)	West Midland s (h)	East Midland s (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/ friends (q)	Yes - me/ family/ friends (r)	No (s)	Yes (t)	No (u)
		Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684
An infection	5*	1*	-	4*	-	-	-	-	*	-	-	3	-	1*	4*	-	-	-	5*	3	2*
Diabetes	3*	-	-	3*	-	1*	2	-	*	-	-	-	-	3	*	1	-	1	2	-	3
Other condition/ illness	3*	-	-	3*	1	-	-	-	1	-	-	-	1	1*	2*	1	1	2	1	1	2
Medication/ overuse of medication	2*	-	-	2*	-	-	-	1*	-	1*	-	1*	-	-	2*	-	-	-	2*	-	2*
Cancer	2*	-	-	2*	1	-	-	-	-	-	-	*	1*	-	2*	-	1*	1*	2*	1	1
Blood/ blood disorder/ infection	2*	-	-	2*	-	-	1	-	-	-	-	1	-	1*	1*	-	-	-	2*	1*	1*
High cholesterol	2*	-	-	2*	1	1	-	-	-	-	1	-	-	-	2*	-	-	-	2*	1*	1*
Sugar	1*	1	-	-	-	-	-	-	-	-	-	-	-	1*	-	-	-	-	1*	-	1*
Genetics/ family history	1*	-	-	1*	-	-	*	-	-	1	-	-	-	-	1	-	-	-	1*	-	1*
Other answers	5*	-	-	5*	-	-	1	-	1	1	2	1	-	1	4	-	2	2	4	2	3
None/ Don't know	409	35	23	351	26	47	36	34	22	36	69	47	33	37	346	12	22	34	369	30	367
	20%	19%	23%	20%	31%afgH20%IJLM	20%	21%	19%	15%	19%	25%IL	17%	19%	15%	20%n	20%Qr	8%	10%	22%QR	14%	21%T

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

* small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 35
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
Second mention

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
A poor diet/ eating bad food (non-specific/ other)	189 9%	94 9%	96 9%	17 8%	34 10%	52 16%	35 10%	18 6%	34 7%	51 9%	87 13%	51 7%	62 11%	53 9%	47 11%	27 5%	29 11%	99 11%
Smoking/ cigarettes	169 8%	91 9%	78 8%	13 6%	22 6%	24 7%	37 10%	30 10%	42 9%	35 6%	62 9%	72 9%	45 8%	42 7%	36 9%	48 9%	27 10%	71 8%
Fatty food	159 8%	60 6%	99 10%A	7 3%	19 5%	23 7%	26 7%	40 13%	45 10%	25 4%	49 7%	85 11%	57 11%	37 6%	32 8%	33 7%	21 8%	63 7%
Drug use/ abuse	156 8%	69 7%	87 8%	20 9%	30 9%	26 8%	29 8%	14 5%	38 8%	50 9%	55 8%	52 7%	43 8%	44 8%	35 9%	34 7%	27 10%	63 7%
NET: Any mentions of Alcohol	94 5%	46 5%	48 5%	14 6%	13 4%	17 5%	13 4%	14 5%	22 5%	27 5%	30 4%	37 5%	27 5%	26 5%	20 5%	21 4%	17 6%	32 4%
Alcohol/ drinking alcohol	67 3%	31 3%	36 3%	13 6%	9 3%	12 4%	8 2%	10 3%	15 3%	22 4%	20 3%	25 3%	17 3%	23 4%	16 4%	12 2%	14 5%	21 2%
Excessive alcohol consumption	17 1%	11 1%	6 1%	- -	2 1%	2 1%	3 1%	3 1%	7 1%	2 *	5 1%	10 1%	7 1%	1 *	3 1%	6 1%	1 *	8 1%
Alcoholism/ alcohol addiction/ abuse	9 *	4 *	5 1%	1 *	2 1%	2 1%	2 1%	2 1%	- -	2 *	4 1%	2 *	3 *	2 *	1 *	3 1%	2 1%	3 *
Obesity/ being overweight	91 5%	32 3%	59 6%A	6 3%	19 5%	11 3%	17 5%	21 7%	17 4%	25 4%	28 4%	39 5%	25 5%	20 3%	14 3%	32 6%	16 6%	34 4%
Hepatitis	68 3%	31 3%	37 4%	4 2%	7 2%	8 2%	19 5%	10 3%	21 4%	11 2%	26 4%	31 4%	17 3%	20 4%	17 4%	14 3%	14 5%	25 3%
Medication/ overuse of medication	36 2%	18 2%	18 2%	6 3%	12 3%	3 1%	7 2%	1 *	7 1%	18 3%	11 2%	7 1%	7 1%	12 2%	10 2%	8 2%	9 3%	14 2%
Genetics/ family history	33 2%	13 1%	20 2%	6 3%	4 1%	5 2%	3 1%	7 2%	8 2%	10 2%	8 1%	15 2%	7 1%	11 2%	5 1%	10 2%	2 1%	14 2%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 35

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Second mention

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Diabetes	25 1%	12 1%	13 1%	1 1%	- -	1 *	2 1%	5 2%D	15 3% cDEF	1 *	4 1%	19 3% J	7 1%	7 1%	6 2%	4 1%	2 1%	3 *
Cancer	23 1%	12 1%	11 1%	- -	- -	1 *	2 1%	4 1% d	14 3% CDEF	- -	4 1%	19 2% J	3 1%	5 1%	6 1%	9 2%	- -	2 *
Sugar	22 1%	11 1%	12 1%	6 3% dEf	2 1%	1 *	3 1%	5 2% e	5 1%	9 2%	3 *	11 1% j	7 1%	4 1%	8 2%	4 1%	2 1%	11 1%
Dehydration/ lack of water/ fluids	17 1%	7 1%	10 1%	1 *	7 2% gh	3 1%	4 1%	* *	2 *	8 1% k	7 1%	2 *	2 *	8 1% l	3 1%	4 1%	6 2%	8 1%
Unhealthy lifestyle (non-specific)	15 1%	3 *	12 1% A	- -	5 1%	4 1%	1 *	1 *	4 1%	5 1%	5 1%	5 1%	7 1%	3 *	1 *	5 1%	1 *	7 1%
An infection	13 1%	10 1% B	2 *	1 1%	1 *	5 1% g	2 *	- -	4 1%	3 *	6 1%	4 *	3 1%	7 1%	1 *	2 *	2 1%	4 *
Other condition/ illness	11 1%	2 *	8 1%	- -	1 *	- -	1 *	3 1% e	5 1% e	1 *	1 *	8 1% J	2 *	6 1% o	2 1%	1 *	2 1%	2 *
Salt	10 *	5 *	5 *	2 1%	3 1%	2 1%	- -	1 *	2 *	5 1%	2 *	2 *	4 1%	1 *	2 *	3 1%	2 1%	3 *
Blood/ blood disorder/ infection	10 *	8 1% b	2 *	2 1%	- -	- -	2 *	3 1%	3 1%	2 *	2 *	6 1%	- -	3 1%	4 1% l	3 1% l	- -	4 *
Gluttony/ overeating	8 *	8 1% B	- -	3 1% eF	2 1%	- -	- -	2 1%	2 *	5 1% J	- -	3 *	4 1% mo	- -	5 1% MO	- -	- -	7 1%
Lack of exercise	8 *	6 1%	2 *	1 *	4 1%	- -	1 *	1 *	2 *	5 1% j	1 *	3 *	3 1%	1 *	1 *	4 1%	- -	7 1%
A virus	6 *	3 *	3 *	2 1% f	- -	2 1%	- -	1 *	1 *	2 *	2 *	1 *	1 *	1 *	1 *	3 1%	- -	2 *
High cholesterol	4 *	- -	4 * a	- -	- -	- -	1 *	1 *	1 *	- -	1 *	2 *	2 *	1 *	1 *	- -	- -	1 *

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 35

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Second mention

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Too much meat/ red meat	1 *	- -	1 *	- -	- -	- -	- -	1 *	- -	- -	- -	1 *	- -	- -	- -	1 *	- -	1 *
Other answers	26 1%	13 1%	12 1%	2 1%	8 2%	3 1%	4 1%	4 1%	5 1%	9 2%	7 1%	9 1%	5 1%	12 2%	4 1%	5 1%	9 3%Q	7 1%
None/ Don't know	843 42%	441 45%B	402 39%	117 52% EFGH	158 45% gH	136 42%	151 42%	110 37%	170 37%	275 48% jK	287 42% k	281 37%	206 38%	246 44% l	157 38%	233 46% LN	93 33%	406 46% P

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 36

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Second mention

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humber (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
A poor diet/ eating bad food (non-specific/ other)	189 9%	12 7%	12 12%	165 9%	7 8%	25 11%	12 7%	20 11%	16 10%	19 10%	28 10%	26 9%	14 8%	26 10%	161 10%	6 9%	21 8%	27 8%	163 10%	20 9%	168 9%
Smoking/ cigarettes	169 8%	19 11%h	5 5%	145 8%	9 10%	21 9%	18 11%h	9 5%	10 7%	14 7%	20 8%	25 9%	18 10%	23 9%	141 8%	3 5%	22 8%	26 8%	144 9%	14 7%	152 9%
Fatty food	159 8%	9 5%	9 9%	141 8%	4 5%	23 10%G	7 4%	11 6%	15 10%g	18 10%g	21 8%	21 8%	20 11%AG	23 9%	133 8%	7 11%	24 9%	29 9%	128 8%	26 12%U	129 7%
Drug use/ abuse	156 8%	21 12%dfgIK	6 6%	129 7%	6 7%	15 6%	10 6%	20 11%lk	4 3%	19 10%lk	13 5%	27 9%l	15 8%i	17 7%	132 8%	1 1%	33 12%PS	33 10%Ps	122 7%	13 6%	142 8%
NET: Any mentions of Alcohol	94 5%	7 4%	2 2%	85 5%	3 4%	7 3%	10 6%	10 5%	11 8%bfK	9 4%	7 3%	19 7%fk	8 5%	13 5%	75 4%	1 1%	20 7%S	20 6%	72 4%	11 5%	82 5%
Alcohol/ drinking alcohol	67 3%	4 2%	2 2%	62 4%	3 4%	6 3%	8 5%K	10 5%K	7 4%k	6 3%	3 1%	14 5%K	5 3%	9 4%	55 3%	-	12 4%	12 4%	55 3%	7 3%	60 3%
Excessive alcohol consumption	17 1%	2 1%	1 1%	15 1%	-	-	2 1%	-	4 2%Fh	2 1%	2 1%	5 2%f	1 *	2 1%	15 1%	-	6 2%S	6 2%S	11 1%	3 1%	15 1%
Alcoholism/ alcohol addiction/ abuse	9 *	1 1%	-	8 *	-	1 1%	-	-	1 1%	1 *	2 1%	1 *	2 1%	2 1%	6 1%	1 1%	2 1%	3 1%	6 *	1 1%	7 *
Obesity/ being overweight	91 5%	5 3%	3 3%	83 5%	4 4%	8 3%	12 7%af	9 5%	9 6%	9 5%	10 4%	13 5%	8 5%	15 6%	74 4%	5 8%	25 9%S	29 9%S	62 4%	22 10%U	68 4%
Hepatitis	68 3%	6 4%	4 4%	57 3%	1 1%	8 3%	8 5%	9 5%j	9 6%eJ	3 2%	6 2%	8 3%	5 3%	13 5%o	51 3%	7 11%S	13 5%	18 6%S	50 3%	10 5%	57 3%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 36

Q3. Can you name three things that you think would make you more likely to develop liver disease?**Base: All respondents****Second mention**

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease				Aware of British Liver Trust	
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/ friends (q)	Yes - me/ family/ friends (r)	No (s)	Yes (t)	No (u)
		Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684
Medication/ overuse of medication	36 2%	2 1%	1 1%	32 2%	2 2%	5 2%	4 2%	3 2%	5 3%K	3 1%	1 *	5 2%	4 2%	8 3%o	28 2%	3 6%S	11 4%S	14 4%S	22 1%	7 3%	29 2%
Genetics/ family history	33 2%	1 *	1 1%	32 2%	1 2%	5 2%	2 1%	2 1%	5 3%al	6 3%a	7 3%	2 1%	1 1%	3 1%	28 2%	-	2 1%	2 1%	29 2%	6 3%	25 1%
Diabetes	25 1%	3 2%	1 1%	21 1%	1 1%	2 1%	1 1%	2 1%	3 2%	3 2%	2 1%	3 1%	3 2%	5 2%	19 1%	2 3%	4 1%	5 2%	18 1%	6 3%u	19 1%
Cancer	23 1%	1 1%	2 2%k	20 1%	3 4%a	1 FK *	1 1%	2 1%	4 2%fk	2 1%	-	4 2%k	2 1%	2 1%	20 1%	-	8 3%S	8 2%S	15 1%	5 2%	18 1%
Sugar	22 1%	4 2%Fk	-	19 1%	-	-	* *	3 2%fk	1 *	9 4%BeFGI-KI	-	4 1%	2 1%f	5 2%	16 1%	-	3 1%	3 1%	18 1%	-	21 1%
Dehydration/ lack of water/ fluids	17 1%	5 3%DJ	-	12 1%	-	1 1%	2 1%	1 *	2 1%	-	3 1%	2 1%	2 1%	-	16 1%	-	3 1%	3 1%	14 1%	4 2%	13 1%
Unhealthy lifestyle (non-specific)	15 1%	-	1 1%f	14 1%	-	-	4 3%AFilm *	1 *	1 1%	* *	5 2%f	1 1%	1 *	* *	15 1%	1 1%	-	1 *	14 1%	1 *	14 1%
An infection	13 1%	-	-	13 1%	2 2%afhi	1 *	1 *	-	-	2 1%	1 *	2 1%	4 2%afhi	-	13 1%	-	4 1%	4 1%	9 1%	1 1%	11 1%
Other condition/ illness	11 1%	4 2%Dikl	1 1%	6 *	-	1 *	1 *	* *	-	1 1%	1 *	1 *	2 1%	2 1%	9 1%	1 1%	3 1%	3 1%	8 *	1 *	10 1%
Salt	10 *	-	-	10 1%	-	1 *	1 1%	-	3 2%L	1 1%	3 1%	-	1 1%	2 1%	8 *	1 1%	4 2%S	4 1%S	6 *	2 1%	8 *
Blood/ blood disorder/ infection	10 *	1 1%	2 2%df	7 *	-	-	-	1 *	-	2 1%	2 1%	1 *	2 1%	1 *	8 *	-	2 1%	2 1%	7 *	3 2%U	6 *

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 36

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Second mention

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotlan d (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshi re & Humbers ide (g)	West Midland s (h)	East Midland s (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/ friends (q)	Yes - me/ family/ friends (r)	No (s)	Yes (t)	No (u)
		Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684
Gluttony/ overeating	8 *	1 1%	-	7 *	-	-	-	-	-	-	3 1%	3 1%	1 *	1 *	8 *	-	-	-	8 *	3 2%U	5 *
Lack of exercise	8 *	-	-	8 *	-	-	-	2 1%	1 *	-	1 *	3 1%	2 1%	1 *	7 *	-	4 2%S	4 1%S	4 *	1 *	7 *
A virus	6 *	-	1 1%dl	5 *	-	-	-	-	1 *	1 *	2 1%	-	1 *	1 1%	4 *	1 1%	-	1 *	5 *	3 1%U	3 *
High cholesterol	4 *	-	-	4 *	-	*	-	1 1%	1 1%	-	-	1 *	-	*	3 *	-	-	-	4 *	-	4 *
Too much meat/ red meat	1 *	-	-	1 *	-	-	-	-	1 1%	-	-	-	-	-	1 *	-	-	-	1 *	-	1 *
Other answers	26 1%	1 *	1 1%	24 1%	1 2%	5 2%	2 1%	3 2%	1 *	2 1%	3 1%	6 2%	1 1%	8 3%O	18 1%	4 6%S	5 2%	8 3% _s	18 1%	6 3%	20 1%
None/ Don't know	843 42%	78 44%i	48 47%llm	717 41%	41 49%llm	104 45%lm	74 43%i	75 41%	49 33%	73 37%	134 50%lJLM	104 37%	63 35%	82 33%	719 42%N	22 36%q	61 23%	81 25%	752 45%QR	57 26%	771 43%T

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 37

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Third mention

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
A poor diet/ eating bad food (non-specific/ other)	124 6%	54 6%	69 7%	16 7%G	25 7%G	24 7%G	19 5%	8 3%	32 7%G	40 7%	43 6%	40 5%	38 7%	35 6%	24 6%	27 5%	22 8%	46 5%
Obesity/ being overweight	92 5%	40 4%	52 5%	12 5%e	19 5%e	7 2%	12 3%	16 5%e	27 6%E	31 5%j	19 3%	42 6%J	24 5%	28 5%	19 5%	20 4%	13 5%	39 4%
Drug use/ abuse	87 4%	44 4%	43 4%	12 5%	14 4%	12 4%	21 6%H	15 5%	14 3%	25 4%	33 5%	29 4%	25 5%	20 4%	18 4%	23 5%	23 8%Q	32 4%
Smoking/ cigarettes	75 4%	34 3%	42 4%	7 3%	13 4%	11 3%	8 2%	13 4%	23 5%F	20 4%	19 3%	36 5%j	24 4%O	19 3%	24 6%O	9 2%	9 3%	26 3%
Genetics/ family history	48 2%	11 1%	37 4%A	7 3%	9 3%	10 3%	7 2%	9 3%	6 1%	16 3%	17 2%	15 2%	15 3%	11 2%	7 2%	14 3%	6 2%	22 2%
Hepatitis	35 2%	18 2%	17 2%	- -	4 1%	7 2%c	6 2%c	10 3%Ch	6 1%	4 1%	13 2%	17 2%i	12 2%	8 1%	6 1%	9 2%	4 2%	10 1%
Fatty food	34 2%	15 2%	19 2%	5 2%	6 2%	3 1%	10 3%g	2 1%	9 2%	10 2%	13 2%	11 1%	8 2%	11 2%	10 2%	6 1%	5 2%	16 2%
NET: Any mentions of Alcohol	32 2%	15 2%	17 2%	1 *	4 1%	6 2%	7 2%	5 2%	9 2%	5 1%	13 2%	14 2%	7 1%	8 1%	13 3%O	4 1%	1 *	17 2%
Alcohol/ drinking alcohol	27 1%	14 1%	14 1%	1 *	4 1%	3 1%	7 2%	4 1%	9 2%	5 1%	10 1%	13 2%	7 1%	6 1%	10 3%o	4 1%	1 *	13 2%
Excessive alcohol consumption	3 *	1 *	2 *	- -	- -	2 1%h	- -	1 *	- -	- -	2 *	1 *	- -	1 *	2 1%	- -	- -	3 *
Alcoholism/ alcohol addiction/ abuse	2 *	1 *	1 *	- -	- -	1 *	- -	- -	1 *	- -	1 *	1 *	1 *	1 *	- -	- -	- -	- -
Lack of exercise	31 2%	20 2%	11 1%	2 1%	6 2%	6 2%	4 1%	4 1%	9 2%	8 1%	10 1%	13 2%	10 2%	8 1%	8 2%	5 1%	2 1%	15 2%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 37

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Third mention

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Diabetes	30 1%	12 1%	19 2%	2 1%	2 1%	3 1%	7 2%	9 3% ^d	8 2%	4 1%	10 1%	16 2% ⁱ	10 2%	12 2%	3 1%	5 1%	8 3% ^q	9 1%
Other condition/ illness	28 1%	14 1%	15 1%	1 *	4 1%	6 2%	7 2%	3 1%	8 2%	5 1%	13 2%	11 1%	8 1%	5 1%	9 2%	8 1%	7 3%	9 1%
Cancer	27 1%	15 1%	12 1%	-	2 1%	4 1%	5 1%	5 2% ^c	12 2% ^{Cd}	2 *	9 1%	16 2% ^l	10 2% ^O	10 2% ^O	6 1%	2 *	4 1%	13 1%
Sugar	15 1%	8 1%	7 1%	1 1%	2 1%	5 2% ^f	1 *	1 *	5 1%	3 1%	6 1%	6 1%	10 2% ^{MO}	-	4 1% ^M	1 *	2 1%	8 1%
Medication/ overuse of medication	13 1%	7 1%	7 1%	1 *	1 *	3 1%	2 1%	4 1%	3 1%	1 *	5 1%	7 1%	3 1%	6 1%	2 1%	2 *	1 *	3 *
An infection	13 1%	6 1%	7 1%	1 1%	1 *	3 1%	2 *	3 1%	3 1%	2 *	5 1%	6 1%	3 1%	5 1%	1 *	4 1%	3 1%	6 1%
Dehydration/ lack of water/ fluids	10 *	8 1% ^B	2 *	1 *	1 *	-	4 1% ^e	1 *	4 1%	2 *	4 1%	4 1%	2 *	2 *	5 1% ^o	1 *	-	5 1%
Unhealthy lifestyle (non-specific)	7 *	2 *	5 *	-	-	3 1%	1 *	-	3 1%	-	4 1%	3 *	2 *	2 *	1 *	2 *	-	4 *
High cholesterol	7 *	1 *	6 1% ^a	-	1 *	1 *	1 *	2 1%	2 *	1 *	3 *	4 *	3 1%	1 *	1 *	1 *	-	3 *
Blood/ blood disorder/ infection	6 *	3 *	3 *	-	-	* *	1 *	2 1%	3 1%	-	1 *	5 1%	1 *	2 *	-	3 1%	-	1 *
A virus	5 *	2 *	3 *	-	-	2 1%	-	3 1% ^{fH}	-	-	2 *	3 *	3 *	1 *	-	2 *	1 *	3 *
Salt	5 *	4 *	1 *	-	-	1 *	-	1 *	3 1%	-	1 *	4 *	2 *	-	2 1%	1 *	-	3 *
Gluttony/ overeating	3 *	2 *	1 *	2 1%	-	-	-	-	1 *	2 *	-	1 *	1 *	-	1 *	1 *	-	1 *

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 37

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Third mention

	Gender		Age									Social Grade				Employment Sector		
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Other answers	50 2%	35 4%B	15 1%	1 1%	14 4%C	6 2%	8 2%	9 3%c	13 3%c	15 3%	14 2%	21 3%	17 3%	18 3%	6 2%	9 2%	13 5%Q	16 2%
None/ Don't know	1251 62%	620 63%	631 61%	158 70%GH	222 64%	204 63%	226 63%	174 59%	267 57%	380 66%K	430 63%k	441 58%	303 56%	355 63%L	247 59%	346 69%LmN	155 56%	582 66%P

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 38
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
Third mention

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
A poor diet/ eating bad food (non-specific/ other)	124 6%	12 7%	3 3%	109 6%	5 5%	12 5%	8 5%	12 6%	8 5%	11 6%	17 6%	20 7%	16 9%B	9 4%	112 7%	2 3%	16 6%	18 6%	104 6%	12 6%	109 6%
Obesity/ being overweight	92 5%	13 7%hK	3 3%	76 4%	2 2%	13 6%K	10 6%K	5 3%	8 5%k	14 7%hK	4 2%	13 4%	7 4%	14 6%	75 4%	3 5%	19 7% _s	22 7% _s	70 4%	17 8% _U	75 4%
Drug use/ abuse	87 4%	4 2%	3 3%	80 5%	5 6%	8 3%	5 3%	11 6% _a	8 6%	6 3%	18 7% _a	10 4%	8 5%	13 5%	74 4%	5 8%	15 6%	20 6% _s	65 4%	15 7% _u	70 4%
Smoking/ cigarettes	75 4%	6 4%	2 2%	66 4%	1 1%	9 4%	13 7% _{beh} lK	5 3%	3 2%	7 4%	4 1%	17 6% _{ei} K	8 4%	10 4%	64 4%	5 8% _s	13 5%	17 5%	57 3%	10 5%	64 4%
Genetics/ family history	48 2%	7 4% _{eg}	4 4% _{eg}	37 2%	- -	6 3%	1 1%	6 3%	5 3%	3 1%	5 2%	4 1%	8 4% _e l	6 3%	40 2%	1 2%	7 3%	8 2%	40 2%	9 4% _u	38 2%
Hepatitis	35 2%	3 2%	3 3% _h	29 2%	3 4% _H k	4 2%	4 2%	1 *	3 2%	5 2%	2 1%	5 2%	3 2%	8 3% _o	27 2%	2 3%	5 2%	7 2%	28 2%	14 6% _U	21 1%
Fatty food	34 2%	1 *	- -	33 2%	1 1%	3 1%	4 2%	8 4% _{AB} fJM	3 2%	2 1%	4 2%	7 3%	1 *	2 1%	30 2%	- -	8 3%	8 2%	26 2%	4 2%	29 2%
NET: Any mentions of Alcohol	32 2%	- -	3 3% _{AL}	28 2%	- -	2 1%	2 1%	3 1%	7 5% _{Ae} FgJ Lm	2 1%	10 4% _{AL}	2 1%	2 1%	5 2%	27 2%	3 5% _s	5 2%	8 2%	24 1%	5 2%	27 1%
Alcohol/ drinking alcohol	27 1%	- -	1 1%	26 1%	- -	2 1%	2 1%	1 *	7 5% _{Ae} FgH JLM	2 1%	10 4% _{Ah} Lm	2 1%	1 1%	3 1%	24 1%	2 4%	4 1%	6 2%	21 1%	4 2%	22 1%
Excessive alcohol consumption	3 *	- -	1 1% _{DI}	2 *	- -	- -	- -	2 1%	- -	- -	- -	- -	- -	- -	3 *	- -	1 *	1 *	2 *	- -	3 *

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
*** small base; ** very small base (under 30) ineligible for sig testing**



Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 38
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
Third mention

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humber (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)	
		Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217
Alcoholism/ alcohol addiction/ abuse	2*	-	1% D	1*	-	-	-	-	-	-	-	-	1*	2% O	-	1% S	-	1*	1*	1*	1*	1*
Lack of exercise	31 2%	2 1%	2 2%	26 2%	4 4%	1 FgJ1%	1 1%	3 2%	6 4%	1 FgjM1%	5 1%	4 1%	1 1%	5 2%	26 2%	-	7 3%	7 2%	24 1%	5 2%	26 1%	
Diabetes	30 1%	3 2%	1 1%	26 2%	1 2%	3 1%	2 1%	3 1%	-	3 2%	1*	9 3% k	4 2% i	8 3% O	21 1%	-	8 3% s	8 2%	22 1%	7 3% u	23 1%	
Other condition/ illness	28 1%	1 1%	1 1%	26 1%	1 1%	1 1%	-	2 1%	2 1%	2 1%	6 2% g	5 2% g	6 3% FG	11 5% O	17 1%	2 4%	6 2%	6 2%	22 1%	10 4% U	19 1%	
Cancer	27 1%	1 1%	* 1%	25 1%	1 2%	3 1%	1*	* 1%	1*	5 3% h	3 1%	5 2%	6 4% aGH12%	6 2%	20 1%	1 1%	5 2%	5 2%	22 1%	3 1%	24 1%	
Sugar	15 1%	1*	2% hm	12 1%	-	4 2%	1*	-	1 1%	1*	1*	5 2%	-	3 1%	13 1%	-	5 2% S	5 2% s	10 1%	1*	14 1%	
Medication/ overuse of medication	13 1%	1 1%	-	12 1%	* 1%	5 2% lm	1 1%	1*	1 1%	2 1%	1*	1*	-	4 2% O	8*	2 3%	2 1%	3 1%	10 1%	1 1%	12 1%	
An infection	13 1%	2 1%	1 1%	11 1%	1 2% f	-	2 1%	3 2% f	3 2% f	1*	* 1%	* 1%	-	2 1%	11 1%	1 2%	2 1%	3 1%	10 1%	3 1%	10 1%	
Dehydration/ lack of water/ fluids	10*	-	-	10 1%	-	-	-	-	4 3% AFGHK1%	-	-	3 1%	1 1%	2 1%	8*	-	3 1%	3 1%	7*	-	10 1%	
Unhealthy lifestyle (non-specific)	7*	-	-	7*	-	2 1%	* 1%	1*	1*	1 1%	1*	1*	-	-	5*	-	1*	1*	6*	1*	6*	
High cholesterol	7*	2 1%	-	5*	-	-	-	1*	-	1*	1 1%	2 1%	1*	1*	6*	-	2 1%	2 1%	5*	2 1%	5*	
Blood/ blood disorder/ infection	6*	-	-	6*	-	2 1%	-	2 1%	1 1%	1*	-	1*	-	2 1%	4*	-	1*	1*	5*	1 1%	5*	

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 38
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
Third mention

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
		Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684
A virus	5*	1*	1%	4*	-	-	-	-	-	1%	2%	-	1*	1%	4*	-	1*	1*	4*	1%	3*
Salt	5*	1*	-	4*	-	-	1%	-	-	-	-	3%	-	-	5*	-	-	-	5*	-	5*
Gluttony/ overeating	3*	1*	1%	2*	-	-	-	-	1*	-	1*	-	-	-	2*	-	1*	1*	2*	2%	1%
Other answers	50%	2%	1%	47%	4%	3%	6%	9%	3%	5%	3%	7%	7%	9%	40%	3%	9%	13%	36%	9%	39%
None/ Don't know	1251	115	70	1066	55	150	109	110	84	119	180	159	102	128	1061	31	127	156	1083	86	1145
	62%	65%	69%	61%	64%	64%	64%	60%	56%	61%	67%	56%	57%	51%	63%	51%	47%	48%	64%	40%	64%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 39

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Top 3 mentions

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
NET: Any mentions of Alcohol	1578 78%	755 76%	823 79%	147 65%	263 76% ^C	242 74% ^C	291 81% ^{Ce}	252 85% ^{CDE}	383 82% ^{CdE}	411 71%	533 78% ^l	635 83% ^{lJ}	436 80% ^O	442 78% ^o	334 80% ^O	367 73%	218 78%	670 75%
Alcohol/ drinking alcohol	1094 54%	509 52%	585 56% ^A	101 44%	190 55% ^c	160 49%	204 57% ^{Ce}	179 60% ^{CE}	260 56% ^{Ce}	290 51%	364 53%	439 57% ^l	289 53%	308 55%	240 58% ^o	256 51%	162 58%	475 54%
Excessive alcohol consumption	346 17%	183 19%	162 16%	28 13%	42 12%	56 17% ^d	57 16%	54 18% ^d	109 23% ^{CDeF}	70 12%	113 17% ⁱ	163 21% ^{lJ}	100 18%	100 18%	68 16%	78 15%	28 10%	135 15% ^p
Alcoholism/ alcohol addiction/ abuse	141 7%	65 7%	76 7%	18 8% ^H	32 9% ^H	26 8% ^H	30 8% ^H	20 7% ^H	14 3%	50 9% ^K	56 8% ^K	35 5%	47 9%	34 6%	27 7%	33 7%	29 10%	61 7%
A poor diet/ eating bad food (non-specific/ other)	336 17%	163 17%	173 17%	33 14% ^g	67 19% ^G	81 25% ^{CFGH}	54 15% ^G	26 9%	76 16% ^G	100 17% ^k	135 20% ^K	102 13%	105 19% ^O	93 16% ^o	79 19% ^O	60 12%	53 19%	155 17%
Smoking/ cigarettes	290 14%	145 15%	145 14%	34 15%	42 12%	44 14%	48 13%	48 16%	74 16%	76 13%	93 13%	122 16%	75 14%	77 14%	72 17%	67 13%	44 16%	111 13%
Drug use/ abuse	248 12%	115 12%	133 13%	32 14%	45 13%	39 12%	51 14%	30 10%	52 11%	76 13%	89 13%	82 11%	70 13%	64 11%	56 13%	58 12%	50 18% ^Q	98 11%
Fatty food	205 10%	80 8%	125 12% ^A	13 6%	28 8%	29 9%	38 11% ^c	43 15% ^{CDE}	54 12% ^C	40 7%	67 10%	98 13% ^{lj}	71 13% ^{MO}	50 9%	43 10%	41 8%	29 10%	84 9%
Obesity/ being overweight	192 9%	75 8%	117 11% ^A	19 8%	39 11% ^e	20 6%	32 9%	37 13% ^E	45 10%	58 10%	52 8%	82 11% ^j	55 10%	48 9%	33 8%	55 11%	31 11%	77 9%
Hepatitis	114 6%	53 5%	61 6%	4 2%	12 3%	17 5% ^c	25 7% ^{Cd}	23 8% ^{CD}	33 7% ^{Cd}	16 3%	43 6% ^l	56 7% ^l	31 6%	33 6%	25 6%	26 5%	20 7% ^q	36 4%
Genetics/ family history	82 4%	26 3%	57 5% ^A	12 6%	13 4%	15 5%	10 3%	16 6%	15 3%	26 4%	25 4%	31 4%	23 4%	23 4%	13 3%	23 5%	8 3%	37 4%
Diabetes	58 3%	24 2%	34 3%	3 1%	2 1%	6 2%	10 3% ^D	14 5% ^{cDe}	23 5% ^{CDE}	5 1%	16 2% ⁱ	37 5% ^{lJ}	18 3%	20 4%	9 2%	10 2%	13 4% ^Q	13 1%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 39

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Top 3 mentions

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Cancer	52 3%	29 3%	23 2%	* *	2 1%	5 1%	8 2% ^c	10 3% ^{CD}	27 6% ^{CDEF}	2 *	12 2% ⁱ	37 5% ^J	13 2%	15 3%	14 3%	10 2%	4 1%	17 2%
Medication/ overuse of medication	52 3%	27 3%	25 2%	8 3%	13 4%	6 2%	10 3%	6 2%	10 2%	20 4%	15 2%	16 2%	10 2%	19 3%	13 3%	10 2%	10 4%	18 2%
Other condition/ illness	42 2%	17 2%	24 2%	1 *	5 2%	6 2%	8 2%	6 2%	15 3% ^C	6 1%	14 2%	21 3% ⁱ	9 2%	12 2%	11 3%	9 2%	8 3%	12 1%
Lack of exercise	39 2%	26 3% ^b	13 1%	2 1%	10 3%	6 2%	5 1%	6 2%	11 2%	13 2%	10 2%	16 2%	12 2%	9 2%	9 2%	9 2%	2 1%	22 2%
Sugar	39 2%	19 2%	20 2%	8 3% ^f	5 2%	6 2%	4 1%	6 2%	10 2%	13 2%	10 1%	16 2%	18 3% ^{MO}	4 1%	12 3% ^{MO}	5 1%	4 1%	20 2%
An infection	30 2%	17 2%	13 1%	5 2%	2 1%	8 2%	4 1%	4 1%	8 2%	7 1%	12 2%	11 2%	10 2%	13 2% ⁿ	2 1%	6 1%	7 3%	12 1%
Dehydration/ lack of water/ fluids	27 1%	15 2%	11 1%	1 1%	8 2% ^g	3 1%	8 2% ^g	1 *	5 1%	9 2%	11 2%	6 1%	4 1%	10 2%	8 2%	5 1%	6 2%	13 1%
Unhealthy lifestyle (non-specific)	22 1%	6 1%	16 2% ^A	- -	5 1%	7 2% ^{cg}	3 1%	1 *	6 1%	5 1%	9 1%	7 1%	9 2%	5 1%	2 *	6 1%	1 *	11 1%
Blood/ blood disorder/ infection	18 1%	12 1%	6 1%	2 1%	- -	* *	3 1%	4 1% ^{de}	8 2% ^{DE}	2 *	3 *	13 2% ^{iJ}	1 *	5 1%	4 1%	8 2% ^L	- -	6 1%
Salt	14 1%	8 1%	6 1%	2 1%	3 1%	3 1% ^f	- -	2 1%	4 1% ^f	5 1%	3 *	6 1%	6 1% ^m	1 *	4 1% ^m	3 1%	2 1%	7 1%
High cholesterol	13 1%	1 *	11 1% ^A	- -	1 *	2 1%	3 1%	3 1%	4 1%	1 *	5 1%	7 1% ⁱ	5 1%	3 *	2 1%	2 *	- -	5 1%
Gluttony/ overeating	12 1%	11 1% ^B	1 *	5 2% ^{EF}	2 1%	- -	- -	2 1%	3 1%	7 1% ^J	- -	5 1% ^j	5 1% ^M	- -	6 1% ^{Mo}	1 *	- -	8 1%
A virus	11 1%	6 1%	5 1%	2 1% ^f	- -	4 1% ^{dfh}	- -	4 1% ^{dFh}	1 *	2 *	4 1%	4 1%	3 1%	1 *	1 *	5 1%	1 *	5 1%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 39

Q3. Can you name three things that you think would make you more likely to develop liver disease?

Base: All respondents

Top 3 mentions

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Too much meat/ red meat	1 *	- -	1 *	- -	- -	- -	- -	1 *	- -	- -	- -	1 *	- -	- -	- -	1 *	- -	1 *
Other answers	76 4%	48 5%B	28 3%	3 1%	20 6%C	10 3%	15 4%	13 4% _c	15 3%	23 4%	25 4%	28 4%	23 4%	29 5% _{no}	11 3%	14 3%	21 7% _Q	28 3%
None/ Don't know	403 20%	212 22%	190 18%	73 32% _{DEFGH}	75 21% _{GH}	74 23% _{GH}	66 18%	43 15%	71 15%	149 26% _{jk}	140 20% _K	114 15%	96 18%	106 19%	75 18%	125 25% _{LMN}	51 18%	200 23%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 40

Q3. Can you name three things that you think would make you more likely to develop liver disease?**Base: All respondents****Top 3 mentions**

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease				Aware of British Liver Trust	
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
NET: Any mentions of Alcohol	1578	143	77	1358	57	184	133	141	127	157	190	227	143	203	1323	45	241	280	1286	175	1383
	78%	80%Ek	77%	78%	67%	79%Ek	78%e	78%e	83%EK	81%EK	70%	81%EK	80%EK	82%	78%	74%	90%PS	87%PS	76%	81%	78%
Alcohol/ drinking alcohol	1094	93	60	941	38	130	101	96	87	116	125	149	98	137	929	29	165	190	894	119	957
	54%	52%	59%ek	54%	45%	56%k	59%EK	52%	58%ek	60%EK	46%	53%	55%	55%	55%	48%	62%S	59%S	53%	55%	54%
Excessive alcohol consumption	346	33	13	300	17	38	21	33	33	28	39	60	31	50	283	9	55	62	282	34	312
	17%	19%	12%	17%	19%	16%	13%	18%	22%bG	14%	15%	21%bGjk	18%	20%	17%	14%	21%	19%	17%	16%	18%
Alcoholism/ alcohol addiction/ abuse	141	17	5	120	2	16	11	14	6	13	25	19	14	17	113	7	20	27	113	22	117
	7%	9%ei	5%	7%	2%	7%	6%	8%e	4%	7%	9%ei	7%	8%e	7%	7%	12%	8%	8%	7%	10%	7%
A poor diet/ eating bad food (non-specific/ other)	336	24	18	293	11	37	20	35	27	29	48	55	32	43	289	10	43	53	282	34	298
	17%	14%	18%	17%	13%	16%	12%	19%g	18%	15%	18%	20%G	18%	17%	17%	16%	16%	16%	17%	16%	17%
Smoking/ cigarettes	290	27	10	254	11	32	34	21	20	23	35	48	28	36	245	11	43	53	236	34	252
	14%	15%	10%	15%	13%	14%	20%BfHJk	12%	13%	12%	13%	17%	16%	15%	14%	18%	16%	17%	14%	16%	14%
Drug use/ abuse	248	27	9	212	11	23	15	32	13	26	30	38	23	30	210	6	50	55	189	28	216
	12%	15%i	9%	12%	13%	10%	9%	18%bFGI	8%	13%	11%	13%	13%	12%	12%	9%	19%S	17%S	11%	13%	12%
Fatty food	205	11	9	185	5	28	12	21	20	23	26	30	22	27	171	7	34	40	162	31	169
	10%	6%	9%	11%a	5%	12%ag	7%	11%	13%Aeg	12%a	10%	11%	12%a	11%	10%	12%	13%	12%	10%	14%u	10%
Obesity/ being overweight	192	18	6	167	6	24	24	14	17	23	14	28	16	29	158	8	44	51	141	43	145
	9%	10%	6%	10%	7%	10%k	14%behK	8%	11%k	12%K	5%	10%k	9%	12%	9%	13%	16%S	16%S	8%	20%U	8%
Hepatitis	114	10	8	95	5	12	11	10	13	8	10	16	9	23	85	8	22	28	86	25	88
	6%	6%	8%	5%	6%	5%	7%	5%	9%k	4%	4%	6%	5%	9%O	5%	14%S	8%S	9%S	5%	11%U	5%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 40
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
Top 3 mentions

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease				Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me/ friends (p)	Yes - my family/ friends (q)	Yes - me/ friends (r)	No (s)	Yes (t)	No (u)
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Genetics/ family history	82 4%	7 4%	5 5%	70 4%	1 2%	11 5%	3 2%	8 4%	10 6%GL	9 5%	13 5%	6 2%	9 5%l	9 4%	69 4%	1 2%	9 3%	10 3%	70 4%	16 7%U	64 4%
Diabetes	58 3%	6 3%	1 1%	50 3%	2 2%	6 3%	5 3%	5 3%	4 2%	6 3%	3 1%	12 4%k	7 4%k	16 6%O	41 2%	3 5%	12 4%	14 4% _s	42 3%	13 6%U	45 3%
Cancer	52 3%	2 1%	2 2%	48 3%	6 7%AFGHK%	3 4%K	2 1%	3 1%	4 3%	7 4%	3 1%	10 4%	9 5%afghK3%	8 3%	43 3%	1 1%	13 5% _S	13 4% _s	39 2%	9 4%	43 2%
Medication/ overuse of medication	52 3%	4 2%	1 1%	47 3%	2 3%	10 4%K	5 3%	4 2%	6 4%K	5 3%	2 1%	7 2%	4 2%	13 5%O	38 2%	5 8% _S	13 5% _S	17 5% _S	35 2%	8 4%	43 2%
Other condition/ illness	42 2%	5 3%g	1 1%	35 2%	2 3%	2 1%	1 *	2 1%	2 2%	4 2%	6 2%	6 2%	9 5%FGH	14 6%O	28 2%	4 6% _s	9 3%	11 3%	31 2%	11 5%U	31 2%
Lack of exercise	39 2%	2 1%	2 2%	34 2%	4 4%FgJ	1 1%	1 1%	5 3%	6 4%FgJ	1 1%	6 2%	7 2%	3 2%	6 2%	33 2%	-	11 4% _S	11 3% _s	28 2%	6 3%	33 2%
Sugar	39 2%	6 3%gK	2 2%	31 2%	-	4 2%	1 1%	3 2%	2 1%	9 5%efGiK*	1 3%k	8 3%k	2 1%	9 3%o	28 2%	-	8 3%	8 2%	30 2%	1 *	37 2%
An infection	30 2%	2 1%	1 1%	27 2%	3 4%FI	1 *	2 1%	3 2%	3 2%	3 2%	4 2%	3 1%	5 3%F	3 1%	27 2%	1 2%	6 2%	7 2%	23 1%	7 3%u	23 1%
Dehydration/ lack of water/ fluids	27 1%	5 3%	-	22 1%	-	1 1%	2 1%	1 *	6 4%beFgH1% jk	2 1%	3 1%	5 2%	3 2%	2 1%	24 1%	-	5 2%	5 2%	21 1%	4 2%	23 1%
Unhealthy lifestyle (non-specific)	22 1%	-	1 1%	21 1%	-	2 1%	5 3%Am	2 1%	2 1%	2 1%	6 2%	2 1%	1 *	* 1%	20 1%	1 1%	1 *	1 *	21 1%	2 1%	20 1%
Blood/ blood disorder/ infection	18 1%	1 1%	2 2%	15 1%	-	2 1%	1 1%	2 1%	1 1%	2 1%	2 1%	2 1%	2 1%	4 2%	13 1%	-	4 1%	4 1%	14 1%	5 3%U	13 1%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 40
Q3. Can you name three things that you think would make you more likely to develop liver disease?
Base: All respondents
Top 3 mentions

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/ friends		No (s)	Yes (t)	No (u)
																	Yes - my family/ friends (q)	Yes - me/ friends (r)			
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Salt	14 1%	1 *	-	14 1%	-	1 *	2 1%	-	3 2%	1 1%	3 1%	3 1%	1 1%	2 1%	13 1%	1 1%	4 2%	4 1%	10 1%	2 1%	13 1%
High cholesterol	13 1%	2 1%	-	11 1%	1 1%	1 *	-	2 1%	1 1%	1 *	2 1%	3 1%	1 *	1 1%	11 1%	-	2 1%	2 1%	11 1%	2 1%	10 1%
Gluttony/ overeating	12 1%	2 1%	1 1%	9 1%	-	-	-	-	1 *	-	4 2%	3 1%	1 *	1 *	10 1%	-	1 *	1 *	11 1%	5 2%U	7 *
A virus	11 1%	1 *	2 2%	9 dfghL *	-	-	-	-	1 *	2 1%	4 2%l	-	1 1%	3 1%	8 *	1 1%	1 *	2 1%	9 1%	4 2%U	6 *
Too much meat/ red meat	1 *	-	-	1 *	-	-	-	-	1 1%	-	-	-	-	-	1 *	-	-	-	1 *	-	1 *
Other answers	76 4%	3 2%	1 1%	72 4%	5 6%ab	8 3%	9 5%b	9 5%b	5 3%	7 4%	8 3%	14 5%	8 4%	14 6%	60 4%	5 8%	16 6% _s	20 6% _S	54 3%	14 7%U	60 3%
None/ Don't know	403 20%	35 19%	23 23%	345 20%	26 31%afgH20% IJLM	47 20%	34 18%	33 15%	22 19%	36 19%	67 25%lL	46 16%	32 18%	36 15%	341 20% _n	12 20% _{Qr}	21 8%	33 10%	365 22% _{QR}	29 13%	362 20% _T

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
* small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 41
Q4. Which of the following, if any, do you think cause or increase the risk of liver disease?
Base: All respondents

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Alcohol	1850 91%	902 91%	949 92%	195 86%	301 86%	297 91%cd	335 93%CD	284 95%CD	439 94%CD	495 86%	632 92%l	723 95%lj	492 91%	511 91%	388 93%	459 92%	247 89%	798 90%
Obesity or being overweight	1166 58%	568 58%	598 58%	123 54%	224 64%cFgh	187 58%	196 55%	168 56%	268 57%	347 60%	384 56%	435 57%	338 62%nO	328 58%	235 56%	265 53%	167 60%	495 56%
Viruses such as Hepatitis B or C	1135 56%	538 54%	597 58%	64 28%	147 42%C	182 56%CD	217 60%CD	203 68%CDEF	321 69%CDEF	211 37%	399 58%l	525 69%lJ	309 57%	310 55%	230 55%	286 57%	147 53%	463 52%
Smoking	815 40%	414 42%	401 39%	106 47%FH	154 44%fH	136 42%h	130 36%	125 42%h	164 35%	260 45%jK	266 39%	289 38%	233 43%	231 41%	157 38%	194 39%	122 44%	368 41%
Type 2 Diabetes	656 32%	337 34%	319 31%	69 31%	104 30%	98 30%	123 34%	107 36%	154 33%	174 30%	221 32%	261 34%	187 35%o	194 34%o	132 32%	143 28%	96 34%	282 32%
Old age	503 25%	282 29%B	221 21%	65 29%	93 27%	88 27%	83 23%	70 24%	105 22%	157 27%	171 25%	175 23%	127 24%	169 30%LnO	99 24%	107 21%	87 31%Q	201 23%
Heart disease	217 11%	103 10%	113 11%	33 15%fH	54 16%FH	37 11%H	33 9%	32 11%H	28 6%	88 15%JK	69 10%	60 8%	73 13%N	59 10%	35 8%	51 10%	34 12%	93 10%
Sun exposure or other forms of radiation exposure	95 5%	52 5%	44 4%	13 6%gH	32 9%eFGH	17 5%gH	15 4%	7 2%	11 2%	45 8%JK	32 5%K	18 2%	36 7%MO	21 4%	21 5%	18 3%	22 8%	45 5%
Pregnancy	58 3%	25 3%	33 3%	10 5%fH	15 4%fH	11 3%h	6 2%	8 3%	7 1%	26 4%K	17 3%	15 2%	21 4%n	18 3%	7 2%	11 2%	19 7%Q	20 2%
Don't know	82 4%	45 5%	36 4%	18 8%fGH	21 6%GH	13 4%	13 4%	6 2%	10 2%	39 7%jK	27 4%k	16 2%	24 4%	31 6%N	10 2%	16 3%	13 5%	43 5%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 42

Q4. Which of the following, if any, do you think cause or increase the risk of liver disease?

Base: All respondents

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland	Wales	NET: England	North East	North West	Yorkshire & Humber	West Midlands	East Midlands	Eastern	London	South East	South West	Yes	No	Yes - me	Yes - my family/friends	Yes - me/family/friends	No	Yes	No
		(a)	(b)	(d)	(e)	(f)	(g)	(h)	(i)	(j)	(k)	(l)	(m)	(n)	(o)	(p)	(q)	(r)	(s)	(t)	(u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Alcohol	1850 91%	168 95% Eh	93 92%	1589 91%	73 86%	218 94% E	159 94% E	162 89%	139 92%	178 92%	244 90%	253 90%	163 92%	226 91%	1563 92%	52 86%	252 94% p	298 92%	1540 91%	201 93%	1624 91%
Obesity or being overweight	1166 58%	100 56%	65 64% Fk	1007 57%	47 55%	118 51%	103 61% f	114 63% Fk	95 62% Fk	108 56%	140 52%	174 62% Fk	102 57%	152 61%	982 58%	31 52%	179 67% pS	206 64% S	951 56%	137 63%	1009 57%
Viruses such as Hepatitis B or C	1135 56%	94 53%	60 60%	981 56%	43 50%	136 58%	103 60%	100 55%	89 58%	103 53%	152 57%	160 57%	96 54%	169 68% O	930 55%	40 66%	192 71% S	226 70% S	901 54%	144 66% U	978 55%
Smoking	815 40%	76 43% fj	33 32%	706 40%	31 36%	79 34%	75 44% bFJ	75 41%	63 42%	64 33%	114 42% j	137 49% BeFJ	68 38% M	82 33%	713 42% N	21 34%	116 43%	134 42%	674 40%	107 49% U	694 39%
Type 2 Diabetes	656 32%	57 32%	39 39% H	559 32%	31 36% h	81 35% h	51 30%	47 26%	52 34%	61 31%	82 31%	89 32%	65 37% H	110 44% O	530 31%	21 35%	103 38% S	121 37% s	529 31%	81 37%	563 32%
Old age	503 25%	45 25%	26 26%	432 25%	23 27%	55 24%	45 27%	49 27%	42 28%	40 21%	52 19%	78 28% k	46 26%	60 24%	430 25%	7 12%	72 27% P	78 24% p	419 25% p	62 28%	435 25%
Heart disease	217 11%	20 11% f	13 13% F	184 11%	12 14% F	13 6%	20 12% F	19 10%	22 15% F	17 9%	29 11% f	35 12% F	17 10%	29 11%	178 10%	8 14%	26 10%	33 10%	178 11%	29 14%	185 10%
Sun exposure or other forms of radiation exposure	95 5%	14 8% DeJ	6 6% j	75 4%	2 2%	11 5%	9 5% j	8 4%	5 3%	4 2%	17 6% j	12 4%	8 5%	11 4%	82 5%	1 2%	15 6%	17 5%	77 5%	15 7%	78 4%
Pregnancy	58 3%	5 3%	3 3%	50 3%	7 8% aFgh	6 12% jLM	4 3%	5 3%	1 *	5 3%	11 4% i	8 3%	4 2%	7 3%	50 3%	1 1%	8 3%	9 3%	48 3%	10 5%	47 3%
Don't know	82 4%	9 5%	6 6% g	67 4%	5 5%	9 4%	3 2%	9 5%	5 3%	5 3%	13 5%	11 4%	8 4%	4 2%	62 4%	1 1%	3 1%	3 1%	75 4% QR	4 2%	72 4%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 43

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Summary table

	Liver disease is almost always caused by alcohol consumption	There are usually no symptoms in the early stages of liver disease	Liver disease is one of the top 5 biggest killers in the UK	Anyone can develop liver disease, regardless of their lifestyle	Bowel cancer causes more deaths than liver disease	Alcohol is more likely to damage the liver if people are overweight	Most people who develop liver disease are alcoholics
Unweighted base	2024	2024	2024	2024	2024	2024	2024
Weighted base	2024	2024	2024	2024	2024	2024	2024
True	901 45%	1513 75%	1617 80%	1669 82%	1128 56%	1549 77%	946 47%
False	1123 55%	511 25%	407 20%	355 18%	896 44%	475 23%	1078 53%

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 44

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Liver disease is almost always caused by alcohol consumption

	Gender		Age									Social Grade				Employment Sector		
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
True	901 45%	450 46%	451 43%	100 44%	149 43%	147 45%	156 43%	117 39%	232 50%fg	249 43%	303 44%	349 46%	223 41%	268 47%l	192 46%	218 44%	109 39%	409 46%p
False	1123 55%	538 54%	586 57%	127 56%	199 57%	179 55%	204 57%h	180 61%H	234 50%	326 57%	382 56%	415 54%	319 59%m	296 53%	225 54%	283 56%	170 61%q	478 54%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 45

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Liver disease is almost always caused by alcohol consumption

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/ friends (q)	Yes - me/ family/ friends (r)	No (s)	Yes (t)	No (u)
																	46%p	36%Q		45%R	44%S
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
True	901 45%	80 45%	48 47%	773 44%	35 41%	99 42%	73 43%	92 50%L	66 43%	81 42%	126 47%	111 39%	91 51%fjL	96 39%	765 45%n	20 33%	97 36%	116 36%	775 46%pQR	97 45%	789 44%
False	1123 55%	98 55%	53 53%	972 56%	50 59%	134 58% m	97 57%	90 50%	86 57%	113 58% m	143 53%	171 61% HM	87 49%	153 61% o	933 55%	40 67% s	171 64% S	207 64% S	909 54%	120 55%	987 56%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 46

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

There are usually no symptoms in the early stages of liver disease

	Gender		Age									Social Grade				Employment Sector		
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
True	1513 75%	712 72%	801 77%A	146 64%	248 71%	245 75%C	277 77%C	222 75%C	374 80%CDg	394 69%	522 76%l	596 78%l	417 77%o	425 75%	311 75%	360 72%	210 75%	652 74%
False	511 25%	275 28%B	236 23%	81 36%EF	100 29%GH	81 25%	83 23%	75 25%h	92 20%	180 31%JK	163 24%	167 22%	125 23%	139 25%	105 25%	142 28%l	68 25%	235 26%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 47

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

There are usually no symptoms in the early stages of liver disease

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
		Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
True	1513	129	76	1308	69	159	123	141	113	152	191	218	142	205	1252	43	223	262	1241	165	1322
	75%	72%	75%	75%	81%F	68%	72%	78%f	74%	78%F	71%	78%F	80%Fk	82%O	74%	72%	83%pS	81%S	74%	76%	74%
False	511	49	26	436	16	74	47	41	39	43	78	63	36	44	445	17	45	61	443	52	454
	25%	28%	25%	25%	19%	32%EhJL	28%	22%	26%	22%	29%m	22%	20%	18%	26%N	28%q	17%	19%	26%QR	24%	26%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 48

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Liver disease is one of the top 5 biggest killers in the UK

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
True	1617 80%	775 78%	843 81%	173 76%	263 76%	266 82% ^d	294 82% ^d	240 81%	381 82% ^d	436 76%	560 82% ^l	621 81% ^l	439 81%	449 80%	335 80%	394 78%	225 81%	695 78%
False	407 20%	213 22%	194 19%	54 24%	85 24% ^{efh}	59 18%	66 18%	57 19%	85 18%	139 24% ^{JK}	125 18%	143 19%	102 19%	115 20%	81 20%	108 22%	54 19%	192 22%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 49

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Liver disease is one of the top 5 biggest killers in the UK

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
True	1617 80%	141 79%	81 80%	1396 80%	68 80%	182 78%	137 81%	151 83%	126 83%	157 81%	206 76%	230 82%	140 78%	197 79%	1354 80%	47 77%	233 87% ^S	274 85% ^S	1327 79%	190 88% ^U	1401 79%
False	407 20%	37 21%	21 20%	349 20%	17 20%	51 22%	33 19%	32 17%	26 17%	37 19%	64 24%	51 18%	39 22%	52 21%	344 20%	14 23%	36 13%	48 15%	357 21% ^{QR}	27 12%	375 21% ^T

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 50

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Anyone can develop liver disease, regardless of their lifestyle

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
True	1669	829	840	177	282	268	303	250	390	458	571	640	448	453	352	417	227	730
	82%	84%	81%	78%	81%	82%	84%	84%	84%	80%	83%	84%	83%	80%	84%	83%	81%	82%
False	355	158	196	50	66	58	57	48	76	116	115	124	93	111	65	85	52	157
	18%	16%	19%	22%	19%	18%	16%	16%	16%	20%	17%	16%	17%	20%	16%	17%	19%	18%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 51

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Anyone can develop liver disease, regardless of their lifestyle

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
		Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
True	1669	156	89	1425	68	198	146	149	128	160	205	228	143	219	1387	53	223	270	1383	187	1463
	82%	88% ^d	88% ^k	82%	80%	85% ^K	86% ^K	82%	84% ^k	82%	76%	81%	80%	88% ^O	82%	88%	83%	84%	82%	86%	82%
False	355	22	12	320	17	35	24	33	24	35	64	53	35	29	311	7	45	52	301	30	313
	18%	12%	12%	18% ^a	20%	15%	14%	18%	16%	18%	24% ^{ABFG}	19% ^a	20% ^a	12%	18% ^N	12%	17%	16%	18%	14%	18%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 52

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Bowel cancer causes more deaths than liver disease

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
True	1128 56%	524 53%	603 58%A	91 40%	198 57%C	193 59%C	198 55%C	168 57%C	280 60%C	288 50%	391 57%I	448 59%I	288 53%	320 57%	236 57%	283 56%	157 56%	500 56%
False	896 44%	463 47%B	433 42%	136 60%DEFGH	150 3%	133 41%	162 45%	129 43%	186 40%	286 50%JK	295 43%	315 41%	253 47%	244 43%	181 43%	219 44%	122 44%	387 44%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 53

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Bowel cancer causes more deaths than liver disease

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
True	1128 56%	97 55%	56 55%	974 56%	56 66% K	127 55% Ghi	89 52% EL	99 54% el	79 52% eL	114 58% EL	135 50% EL	177 63% Ghi	100 56% K	132 53%	954 56%	27 45%	139 52%	161 50% S	957 57% R	105 48% u	1002 56% t
False	896 44%	81 45%	45 45%	770 44%	29 34%	105 45% el	81 48% EL	84 46% el	73 48% eL	81 42%	134 50% EL	104 37%	79 44%	117 47%	743 44%	33 55%	130 48%	161 50% S	727 43%	112 52% u	774 44%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 54

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Alcohol is more likely to damage the liver if people are overweight

	Gender		Age						Social Grade				Employment Sector					
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
True	1549 77%	802 81%B	747 72%	161 71%	280 81%CF	246 76%	257 71%	232 78%f	372 80%CF	441 77%	504 73%	605 79%J	433 80%No	444 79%N	297 71%	376 75%	209 75%	689 78%
False	475 23%	185 19%	290 28%A	66 29%DH	68 19%	79 24%	103 29%DgH	65 22%	94 20%	133 23%	182 27%K	159 21%	109 20%	120 21%	120 29%LM	126 25%l	70 25%	198 22%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 55

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Alcohol is more likely to damage the liver if people are overweight

	Total	Region											GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust			
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
True	1549 77%	145 81%Fl	75 75%	1329 76%	65 76%	167 72%	136 80%fl	138 76%l	101 66%	162 83%bFIL	217 81%fl	210 74%	133 75%	191 77%	1299 77%	45 75%	195 73%	235 73%	1301 77%	163 75%	1363 77%
False	475 23%	33 19%	26 25%j	416 24%	20 24%	66 28%Ag	34 20%Jk	44 24%	51 34%AGh	32 17%	52 19%	72 26%J	45 25%j	58 23%	399 23%	15 25%	73 27%	87 27%	383 23%	54 25%	413 23%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 56

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Most people who develop liver disease are alcoholics

	Gender		Age									Social Grade				Employment Sector		
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
True	946 47%	458 46%	488 47%	131 58%FGH	178 51%fG	165 51%fG	158 44%G	106 36%	208 45%G	309 54%JK	322 47%K	314 41%	227 42%	269 48%l	203 49%l	247 49%L	125 45%	435 49%
False	1078 53%	529 54%	549 53%	96 42%	170 49%	161 49%	202 56%Cde	191 64%CDEFH	258 5%5C	265 46%	363 53%l	449 59%lJ	315 58%mnO	295 52%	213 51%	255 51%	154 55%	452 51%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 57

Q5. For each of the statements below, please indicate whether you think it is true or false. If you're not sure, please provide your best guess.

Base: All respondents

Most people who develop liver disease are alcoholics

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
True	946 47%	83 47%	41 40%	822 47%	35 41%	115 50%	79 46%	89 49%	65 43%	83 43%	142 53%bjl	120 43%	94 53%beij	95 38%	814 48%N	28 46%	106 40%	133 41%	800 48%Qr	107 49%	822 46%
False	1078 53%	95 53%	61 60%km	923 53%	50 59%m	118 50%	91 54%	93 51%	87 57%m	111 57%km	128 47%	161 57%kM	84 47%	154 62%O	883 52%	33 54%	162 60%S	190 59%S	883 52%	110 51%	954 54%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey
ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 58

Q6. Has your GP ever spoken to you about or tested you for any of the following?**Base: All respondents****Summary table**

	Blood pressure levels	Diabetes	Dietary issues	Cholesterol levels	Liver disease	Alcohol consumption
Unweighted base	2024	2024	2024	2024	2024	2024
Weighted base	2024	2024	2024	2024	2024	2024
Yes	1379 68%	710 35%	457 23%	959 47%	249 12%	320 16%
No	615 30%	1264 62%	1518 75%	1016 50%	1697 84%	1675 83%
Don't know	30 2%	50 2%	49 2%	49 2%	78 4%	29 1%

Liver Disease Survey
ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 59

Q6. Has your GP ever spoken to you about or tested you for any of the following?**Base: All respondents****Blood pressure levels**

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Yes	1379 68%	653 66%	726 70% ^a	78 34%	189 54% ^C	196 60% ^C	258 72% ^{CDE}	246 83% ^{CDEF}	412 88% ^{CDEFG}	267 6%	454 66% ^I	658 86% ^{IJ}	363 67%	391 69%	275 66%	350 70%	185 66%	535 60%
No	615 30%	322 33% ^b	293 28%	140 62% ^{DEFGH}	148 3% ^{FGH}	122 38% ^{FGH}	101 28% ^{GH}	52 17% ^H	52 11%	288 50% ^{JK}	223 33% ^K	104 14%	168 31%	166 29%	139 33%	142 28%	94 34%	334 38%
Don't know	30 2%	12 1%	18 2%	9 4% ^{FGH}	11 3% ^{FGH}	7 2% ^{FGH}	1 *	- -	2 *	20 3% ^{JK}	9 1% ^K	2 *	10 2%	8 1%	3 1%	10 2%	- -	18 2% ^P

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 60

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Blood pressure levels

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside	West Midlands	East Midlands	Eastern (i)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends	Yes - me/family/friends	No (s)	Yes (t)	No (u)
							(g)	(h)	(j)								(q)	(r)			
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Yes	1379 68%	124 70%	68 67%	1187 68%	54 64%	155 67%	127 75%efjK	122 67%	105 69%	127 65%	170 63%	199 71%	128 72%	226 91%O	1098 65%	44 73%	213 80%S	251 78%S	1117 66%	152 70%	1206 68%
No	615 30%	51 29%	31 30%	533 31%	31 36%Gm	74 32%g	40 23%	60 33%g	46 31%	63 32%g	96 36%Gm	78 28%	45 25%	23 9%	589 35%N	16 27%	53 20%	69 21%	543 32%QR	65 30%	542 31%
Don't know	30 2%	2 1%	3 3%	25 1%	- -	4 2%	3 2%	1 *	1 *	4 2%	3 1%	4 2%	5 3%h	- -	11 1%	- -	2 1%	2 1%	24 1%	- -	27 2%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 61

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Diabetes

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Yes	710 35%	364 37%	346 33%	32 14%	86 25% ^C	91 28% ^C	131 36% ^{CDE}	150 50% ^{CDEF}	220 47% ^{CDEF}	118 21%	222 32% ^I	370 48% ^{IJ}	186 34%	201 36%	139 33%	183 36%	91 33%	257 29%
No	1264 62%	600 61%	664 64%	186 82% ^{DEFGH}	247 71% ^{fGH}	228 70% ^{fGH}	227 63% ^{GH}	144 49%	231 50%	434 75% ^{JK}	455 66% ^K	375 49%	345 64%	348 62%	269 65%	302 60%	184 66%	609 69%
Don't know	50 2%	23 2%	27 3%	8 4% ^{Fg}	14 4% ^{FG}	6 2%	2 1%	3 1%	16 3% ^{Fg}	23 4% ^J	9 1%	18 2%	11 2%	15 3%	8 2%	17 3%	4 2%	21 2%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 62
Q6. Has your GP ever spoken to you about or tested you for any of the following?
 Base: All respondents
Diabetes

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Yes	710 35%	58 33%	40 40%	612 35%	32 37%	79 34%	58 34%	71 39%	62 41%	68 35%	98 36%	89 32%	55 31%	184 74%	485 29%	36 59%	105 39%	136 42%	567 34%	92 43%	607 34%
No	1264 62%	117 66%	56 56%	1090 63%	48 56%	149 64%	110 64%	105 58%	87 57%	119 61%	165 61%	188 67%	120 67%	58 23%	1195 70%	23 38%	161 60%	183 57%	1074 64%	123 57%	1125 63%
Don't know	50 2%	3 2%	5 4%	42 2%	5 6%	4 2%	3 2%	6 3%	3 2%	8 4%	6 2%	5 2%	3 2%	7 3%	17 1%	2 3%	2 1%	4 1%	43 3%	2 1%	44 2%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 63

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Dietary issues

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Yes	457 23%	239 24%	218 21%	32 14%	54 15%	79 24%CD	79 22%Cd	92 31%CD	121 26%CD	85 15%	158 23%l	213 28%lJ	116 21%	143 25%N	77 18%	121 24%n	63 22%	165 19%
No	1518 75%	722 73%	796 77%a	187 83%EGH	275 79%Gh	240 74%	280 78%Gh	200 67%	336 72%	462 80%jK	520 76%K	537 70%	413 76%	406 72%	335 80%MO	365 73%	211 76%	697 79%
Don't know	49 2%	27 3%	22 2%	8 4%F	20 6%EFGH	7 2%F	1 *	5 2%f	9 2%f	28 5%JK	8 1%	14 2%	13 2%	15 3%	5 1%	16 3%n	5 2%	25 3%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 64

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Dietary issues

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease				Aware of British Liver Trust	
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
		Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Yes	457 23%	31 18%	25 25%M	400 23%	17 21%	54 23%M	34 20%	46 25%M	36 24%M	47 24%M	74 28%AM	67 24%M	26 15%	130 52%O	303 18%	35 58%QRS	67 25%	99 31%S	354 21%	63 29%U	387 22%
No	1518 75%	144 81%dBK	70 69%	1304 75%	64 75%	174 75%	135 79%bK	134 73%	112 74%	142 73%	186 69%	207 74%	151 85%Be	111 5%FH	1375 81%N	26 42%	195 73%P	218 67%P	1292 77%PR	154 71%	1343 76%
Don't know	49 2%	3 2%	6 6%da	41 GhM2%	4 5%gM	5 2%	1 1%	3 2%	3 2%	6 3%	9 3%	7 3%	1 1%	7 3%O	19 1%	-	6 2%	6 2%	38 2%	1 *	46 3%t

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 65

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Cholesterol levels

	Total	Gender		Age									Social Grade				Employment Sector	
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Yes	959 47%	500 51%B	458 44%	22 10%	48 14%	104 32%CD	196 54%CDE	223 75%CDEF	367 79%CDEF	69 12%	300 44%I	590 77%J	245 45%	260 46%	187 45%	267 53%LMN	113 40%	330 37%
No	1016 50%	471 48%	545 53%A	195 86%EF	284 82%FGH	208 64%FGH	162 45%GH	74 25%	94 20%	479 83%JK	369 54%K	168 22%	282 52%O	289 51%O	225 54%O	220 44%	158 57%	535 60%
Don't know	49 2%	17 2%	33 3%a	10 4%FGH	16 5%FGH	14 4%FGH	3 1%	1 *	5 1%	26 5%jK	17 2%K	6 1%	15 3%	15 3%	5 1%	15 3%n	7 3%	22 3%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 66

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Cholesterol levels

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Yes	959 47%	76 43%	54 53%K	829 48%	41 48%	117 50%K	84 50%k	90 49%k	87 57%AK	92 Im47%	105 39%	131 47%	82 46%	203 82%O	708 42%	34 56%	149 55%S	177 55%S	779 46%	107 49%	842 47%
No	1016 50%	99 55%bl	44 43%	873 50%	41 48%	112 48%	83 49%	88 48%	65 43%	96 50%	152 56%BI	146 52%i	91 51%	44 18%	966 57%N	27 44%	118 44%	144 45%	862 51%Qr	101 47%	897 51%
Don't know	49 2%	3 2%	3 3%l	42 2%	4 4%l	3 1%	3 2%	5 3%i	- -	6 3%i	13 5%fl	4 1%	6 3%l	2 1%	23 1%	- -	2 1%	2 1%	43 3%qR	8 4%	37 2%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 67

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Liver disease

	Total	Gender		Age						Social Grade				Employment Sector				
		Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Yes	249 12%	160 16%B	88 9%	9 4%	28 8%	36 11%C	43 12%C	56 19% CDEF	77 16% CDef	36 6%	79 12% I	133 17% J	66 12%	73 13%	47 11%	62 12%	28 10%	88 10%
No	1697 84%	785 79%	912 88%A	210 93% EFGH	313 90% eGH	276 85% GH	308 86% GH	228 77%	362 78%	523 91% JK	584 85% K	590 77%	459 85%	472 84%	349 84%	417 83%	245 88%	772 87%
Don't know	78 4%	42 4%	36 3%	8 4%	7 2%	13 4%	9 3%	13 4%	27 6% DF	16 3%	22 3%	40 5% Ij	16 3%	19 3%	21 5%	22 4%	5 2%	28 3%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 68

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Liver disease

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
	Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Yes	249 12%	18 10%	14 13%	217 12%	11 13%	36 16% _{jL}	24 14% _l	24 13%	23 15% _{jL}	18 9%	30 11%	23 8%	27 15% _{jL}	249 100% _O	-	44 73% _{QRS}	38 14% _s	77 24% _{QS}	172 10%	40 19% _U	206 12%
No	1697 84%	153 86%	82 81%	1462 84%	73 86%	188 81%	143 84%	151 83%	120 79%	164 84%	230 85%	250 89% _{BFHl}	144 81%	-	1697 100% _N	16 27%	221 82% _{PR}	236 73% _P	1447 86% _{PR}	172 79%	1499 84%
Don't know	78 4%	7 4%	6 6% _g	65 4%	1 1%	8 4%	3 2%	8 4%	8 5% _g	13 7% _{eGI}	9 3%	8 3%	7 4%	-	-	-	9 3%	9 3%	65 4%	4 2%	71 4%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 69

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Alcohol consumption

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Yes	320 16%	201 20%B	119 11%	24 10%	46 13%	47 14%	53 15%	62 21% CDEf	88 19% Cd	70 12%	100 15%	150 20% J	97 18% N	104 18% No	47 11%	72 14%	37 13%	131 15%
No	1675 83%	771 78%	904 87%A	196 87% gh	292 84%	274 84%	305 85% gh	235 79%	372 80%	489 85% K	579 84% K	608 80%	434 80%	452 80%	365 88% LM	423 84%	241 87%	738 83%
Don't know	29 1%	15 2%	14 1%	7 3% FG	9 3% FG	5 2% G	2 *	-	6 1% g	16 3% JK	7 1%	6 1%	10 2%	8 1%	4 1%	7 1%	-	18 2% P

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 70

Q6. Has your GP ever spoken to you about or tested you for any of the following?

Base: All respondents

Alcohol consumption

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Yes	320 16%	21 12%	16 16%	282 16%	19 22%AJ	37 16%	26 15%	27 15%	29 19%aj	24 12%	44 16%	47 17%	28 16%	120 48%O	181 11%	28 46%QRS	40 15%	63 20%S	254 15%	34 16%	281 16%
No	1675 83%	154 86%e	81 80%	1440 83%	65 77%	191 82%	144 85%	153 84%	121 80%	166 86%e	221 82%	230 82%	148 83%	127 51%	1511 89%N	33 54%	227 85%P	258 80%P	1405 83%P	183 84%	1469 83%
Don't know	29 1%	3 2%	4 4%dG	23 1%	1 1%	4 2%g	- -	2 1%	1 1%	4 2%g	5 2%	5 2%	1 1%	2 1%	6 *	- -	2 1%	2 1%	24 1%	1 *	26 1%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 71

Q7. Have you or, to your knowledge, any of your family or friends been affected by liver disease of any sort?

Base: All respondents

	Gender		Age									Social Grade				Employment Sector		
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Pri- vate (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
NET: Yes	323 16%	139 14%	184 18%A	26 11%	59 17%	48 15%	46 13%	52 17% _c	92 20% _{CeF}	85 15%	94 14%	144 19% _J	76 14%	89 16%	70 17%	87 17%	44 16%	114 13%
Yes - me	60 3%	33 3%	27 3%	3 1%	16 5% _c	10 3%	9 2%	11 4%	12 3%	19 3%	18 3%	23 3%	10 2%	17 3%	16 4%	17 3%	11 4%	20 2%
Yes - my family or friends	268 13%	108 11%	161 16%A	23 10%	43 12%	40 12%	38 11%	43 14%	81 17% _{CdeF}	66 11%	79 11%	124 16% _J	67 12%	75 13%	54 13%	72 14%	34 12%	96 11%
No	1684 83%	838 85% _b	846 82%	194 86%	282 81%	276 85% _h	313 87% _{dH}	246 83%	373 80%	476 83%	589 86% _K	618 81%	462 85%	471 83%	340 82%	411 82%	232 83%	764 86%
Refused	17 1%	11 1%	6 1%	7 3% _{EFGH}	7 2% _{Gh}	1 *	2 *	- -	1 *	13 2% _{JK}	3 *	1 *	3 1%	4 1%	6 1%	4 1%	2 1%	8 1%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 72

Q7. Have you or, to your knowledge, any of your family or friends been affected by liver disease of any sort?

Base: All respondents

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease				Aware of British Liver Trust	
		Scotlan	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West	Yes	No	Yes - me	Yes - my family/friends	Yes - me/family/friends	No	Yes	No
		(a)	(b)	(d)	(e)	(f)	(g)	(h)	(i)	(j)	(k)	(l)	(m)	(n)	(o)	(p)	(q)	(r)	(s)	(t)	(u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
NET: Yes	323	31	17	274	19	41	24	24	20	32	40	53	20	77	236	60	268	323	-	45	274
	16%	17%	17%	16%	22%hi	18%M	14%	13%	13%	17%	15%	19%M	11%	31%O	14%	100%S	100%S	100%S	-	21%u	15%
Yes - me	60	6	3	51	3	10	4	5	6	4	11	5	2	44	16	60	6	60	-	13	47
	3%	3%	3%	3%	4%	4%M	2%	3%	4%	2%	4%	2%	1%	18%O	1%	100%QRS	2%S	19%QS	-	6%U	3%
Yes - my family or friends	268	25	15	228	15	33	21	19	14	28	31	49	18	38	221	6	268	268	-	34	230
	13%	14%	15%	13%	18%him	14%	12%	10%	9%	14%	11%	17%HIM	10%	15%	13%	10%S	100%PRS	83%PS	-	16%	13%
No	1684	146	84	1454	66	191	145	155	129	159	224	227	158	172	1447	-	-	-	1684	171	1488
	83%	82%	83%	83%	77%	82%	85%	85%	85%	82%	83%	81%	89%EfjL	69%	85%N	-	-	-	100%PQR	79%	84%
Refused	17	1	-	17	1	-	1	3	2	3	5	2	-	-	14	-	-	-	-	1	13
	1%	*	-	1%	1%	-	1%	2%f	2%f	2%f	2%f	1%	-	-	1%	-	-	-	-	1%	1%

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u

*** small base; ** very small base (under 30) ineligible for sig testing**

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 73
Q8. What charity, if any, first comes to mind when you think of liver disease?
Base: All respondents

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
British Liver Trust	138 7%	63 6%	75 7%	35 15% DEFGH	22 6%	15 4%	21 6%	22 7%	24 5%	56 10% JK	36 5%	46 6%	41 8%	30 5%	30 7%	37 7%	12 4%	65 7%
Cancer Research	70 3%	35 4%	35 3%	6 3%	13 4% e	4 1%	11 3%	13 4% E	22 5% E	19 3%	16 2%	35 5% J	17 3%	19 3%	18 4%	16 3%	10 3%	25 3%
Alcoholics Anonymous	35 2%	21 2%	14 1%	1 *	2 1%	4 1%	6 2%	6 2%	17 4% CDEf	3 *	9 1%	23 3% lj	10 2% n	11 2% N	1 *	13 3% N	5 2%	10 1%
Macmillan	29 1%	11 1%	18 2%	3 1%	3 1%	8 2%	3 1%	3 1%	9 2%	6 1%	10 2%	12 2%	4 1%	8 1%	12 3% Lo	5 1%	2 1%	15 2%
British Heart Foundation	20 1%	12 1%	8 1%	1 *	5 1% h	8 2% cFH	1 *	4 1% h	1 *	6 1%	9 1%	5 1%	4 1%	6 1%	7 2%	2 *	9 3% Q	9 1%
Foundation For Liver Research	14 1%	6 1%	8 1%	1 1%	-	2 1%	3 1%	4 1% d	3 1%	1 *	5 1%	7 1%	2 *	2 *	6 1% m	4 1%	-	10 1%
Liver Foundation	13 1%	7 1%	6 1%	-	2 1%	3 1%	3 1%	2 1%	3 1%	2 *	5 1%	5 1%	1 *	1 *	7 2% LM	4 1%	-	8 1%
Cancer UK	12 1%	8 1%	4 *	-	-	5 2% dF	-	4 1% dF	3 1%	-	5 1% i	7 1% l	2 *	4 1%	3 1%	3 1%	2 1%	4 *
Other answers	143 7%	71 7%	72 7%	15 7%	26 8%	20 6%	28 8%	16 5%	39 8%	41 7%	48 7%	55 7%	50 9% no	39 7%	24 6%	30 6%	24 9%	68 8%
None/ Don't know	1577 78%	761 77%	816 79%	166 73%	279 80%	266 82% Cgh	287 80%	225 76%	353 76%	446 78%	553 81% K	578 76%	416 77%	453 80%	314 75%	393 78%	217 78%	689 78%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 74

Q8. What charity, if any, first comes to mind when you think of liver disease?

Base: All respondents

	Region													GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
	Total	Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humberside (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
	Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
British Liver Trust	138	13	5	120	10	15	7	11	9	8	26	18	17	16	120	2	23	25	113	133	4
	7%	7%	5%	7%	11%GJ	6%	4%	6%	6%	4%	10%gj	6%	10%gj	6%	7%	4%	9%	8%	7%	61%U	*
Cancer Research	70	5	1	63	4	5	8	7	8	9	8	11	4	11	58	2	13	15	54	3	67
	3%	3%	1%	4%	5%	2%	5%	4%	5%	4%	3%	4%	2%	4%	3%	3%	5%	5%	3%	1%	4%
Alcoholics Anonymous	35	2	1	32	3	6	2	1	2	5	2	8	4	6	29	2	9	11	24	1	34
	2%	1%	1%	2%	3%h	3%h	1%	*	1%	2%	1%	3%h	2%	2%	2%	3%	3%S	3%S	1%	1%	2%
Macmillan	29	3	3	24	1	1	3	6	2	2	5	3	1	2	26	-	9	9	19	4	25
	1%	1%	3%	1%	1%	*	2%	3%Fm	1%	1%	2%	1%	*	1%	2%	-	4%S	3%S	1%	2%	1%
British Heart Foundation	20	2	-	18	-	3	-	2	*	-	10	2	-	1	19	-	*	*	19	7	13
	1%	1%	-	1%	-	1%	-	1%	*	-	4%bGiJL	1%	-	*	1%	-	*	*	1%	3%U	1%
Foundation For Liver Research	14	-	2	11	-	2	-	1	4	1	2	2	-	5	9	-	1	1	12	8	6
	1%	-	2%dagm	1%	-	1%	-	*	3%AGhM1%	1%	1%	1%	-	2%O	1%	-	1%	*	1%	4%U	*
Liver Foundation	13	-	-	13	1	1	1	3	-	1	3	2	1	2	10	2	4	4	9	8	4
	1%	-	-	1%	1%	*	*	2%a	-	*	1%	1%	1%	1%	1%	3%S	1%	1%	1%	3%U	*
Cancer UK	12	1	1	10	-	2	1	1	2	1	-	3	*	4	8	-	2	2	11	1	11
	1%	*	1%k	1%	-	1%	1%	*	1%	*	-	1%	*	1%o	*	-	1%	1%	1%	1%	1%
Other answers	143	8	9	126	4	11	5	24	7	10	30	22	12	27	108	7	28	35	105	56	80
	7%	5%	9%G	7%	5%	5%	3%	13%AeFGI	5%	5%	11%AFGij	8%G	7%	11%O	6%	12%	10%S	11%S	6%	26%U	4%
None/ Don't know	1577	147	78	1352	63	188	143	132	120	158	195	214	140	177	1335	46	184	225	1339	17	1538
	78%	82%HK	77%	77%	74%	81%hk	84%eHKI	72%	79%	82%hk	73%	76%	78%	71%	79%N	76%	69%	70%	80%QR	8%	87%T

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing



Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 75

Q9. Have you ever heard of the charity the British Liver Trust?**Base: All respondents**

	Gender			Age									Social Grade				Employment Sector	
	Total	Male (a)	Female (b)	18-24 (c)	25-34 (d)	35-44 (e)	45-54 (f)	55-64 (g)	65+ (h)	18-34 (i)	35-54 (j)	55+ (k)	AB (l)	C1 (m)	C2 (n)	DE (o)	Public (p)	Private (q)
Unweighted base	2024	990	1034	216	259	342	392	301	514	475	734	815	566	589	351	518	249	825
Weighted base	2024	988	1036	227	348	326	360	297	466	575	686	764	542	564	416	502	279	887
Yes	217 11%	106 11%	111 11%	42 19%	35 10%	30 9%	40 11%	34 11%	35 8%	78 14%	70 10%	69 9%	63 12%	55 10%	48 11%	50 10%	30 11%	109 12%
No	1776 88%	864 87%	912 88%	182 80%	304 87%	293 90%	313 87%	260 87%	425 91%	486 85%	605 88%	684 90%	467 86%	506 90%	365 88%	438 87%	247 89%	758 86%
Don't know	31 2%	18 2%	13 1%	2 1%	9 3%	3 1%	7 2%	4 1%	6 1%	11 2%	10 1%	10 1%	11 2%	3 *	4 1%	13 3%	1 *	20 2%

Proportions/Means: Columns Tested (5%, 10% risk level) - a/b - c/d/e/f/g/h - i/j/k - l/m/n/o - p/q

Liver Disease Survey

ONLINE Fieldwork: 24th-25th May 2017

Absolutes/col percents

Table 76
Q9. Have you ever heard of the charity the British Liver Trust?
Base: All respondents

	Total	Region												GP has spoken to or tested me for liver disease		Affected by liver disease			Aware of British Liver Trust		
		Scotland (a)	Wales (b)	NET: England (d)	North East (e)	North West (f)	Yorkshire & Humbers (g)	West Midlands (h)	East Midlands (i)	Eastern (j)	London (k)	South East (l)	South West (m)	Yes (n)	No (o)	Yes - me (p)	Yes - my family/friends (q)	Yes - me/family/friends (r)	No (s)	Yes (t)	No (u)
Unweighted base	2024	178	111	1735	94	250	196	188	159	185	191	280	192	266	1679	56	279	328	1680	203	1788
Weighted base	2024	178	101*	1745	85*	233	170	182	152	194	269	281	178	249	1697	60*	268	323	1684	217	1776
Yes	217 11%	16 9%	12 12%G	189 11%	10 12%G	23 10%g	8 5%	21 12%G	13 9%	13 7%	45 17%Af	26 9%G	28 16%af	40 16%Gi	172 10%	13 22%S	34 13%	45 14%S	171 10%	217 100%U	-
No	1776 88%	159 89%km	88 87%	1529 88%	72 84%	209 90%Km	159 94%bEH	153 4%K	139 91%hKM	179 92%eHK	220 82%	252 89%Km	147 83%	206 83%	1499 88%N	47 78%	230 86%	274 85%	1488 88%P	-	1776 100%T
Don't know	31 2%	3 2%	1 1%	27 2%	3 3%fl	2 1%	3 2%	8 4%F	- -	2 1%	3 1%	3 1%	3 1%	2 1%	26 2%	-	4 1%	4 1%	25 1%	-	-

Proportions/Mean: Columns Tested (5%, 10% risk level) - a/d - b/d - c/d - a/b/c/e/f/g/h/i/j/k/l/m - n/o - p/q/r/s - t/u
 * small base; ** very small base (under 30) ineligible for sig testing