

OMNICELL – MEDICATION ADHERENCE

METHODOLOGY NOTE

ComRes interviewed 2,048 British adults online between 31ST July and 2nd August 2015. Data were weighted to be nationally representative of all British adults aged 18+. ComRes is a member of the British Polling Council and abides by its rules. All press releases or other publications must be checked with ComRes before use. ComRes requires 48 hours to check a press release unless otherwise agreed.

To commission a voting intention poll or a public opinion survey please contact Katharine Peacock:
katharine.peacock@comres.co.uk.

To register for Pollwatch, a monthly newsletter update on the polls, please email: pollwatch@comres.co.uk.

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 1
Q.1 When was the last time you were prescribed any type of medicine on the NHS, if ever?
Base: All respondents

	Gender		Age							Social Grade				Employment Sector	
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	2048	1043	1005	202	341	314	379	300	512	581	582	314	571	204	631
Weighted base	2048	997	1051	238	355	336	365	295	459	551	551	446	500	274	885
NET: In the last year	1505 73%	700 70%	805 77%	163 68%	238 67%	217 65%	251 69%	234 79%	402 88%	392 71%	394 71%	332 74%	386 77%	185 67%	591 67%
In the last month	925 45%	464 47%	462 44%	48 20%	110 31%	120 36%	168 46%	166 56%	313 68%	232 42%	225 41%	213 48%	255 51%	102 37%	285 32%
More than 1 month and up to 6 months ago	381 19%	153 15%	228 22%	73 31%	84 24%	55 16%	57 15%	49 17%	64 14%	107 19%	112 20%	74 16%	89 18%	50 18%	198 22%
More than 6 months and up to 1 year ago	198 10%	83 8%	115 11%	42 18%	43 12%	43 13%	26 7%	19 7%	25 5%	53 10%	57 10%	45 10%	43 9%	32 12%	108 12%
More than 1 year and up to 5 years ago	279 14%	143 14%	136 13%	37 15%	60 17%	62 19%	61 17%	28 9%	32 7%	88 16%	86 16%	55 12%	50 10%	44 16%	155 18%
More than 5 years ago	197 10%	117 12%	80 8%	26 11%	31 9%	42 12%	46 13%	29 10%	23 5%	55 10%	46 8%	52 12%	43 9%	34 13%	100 11%
I have never been prescribed any medicine on the NHS	68 3%	38 4%	30 3%	13 5%	26 7%	15 4%	8 2%	4 1%	2 1%	15 3%	25 5%	8 2%	20 4%	12 4%	38 4%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 2

Q.1 When was the last time you were prescribed any type of medicine on the NHS, if ever?

Base: All respondents

	Region												
	Total	Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	2048	174	111	1763	78	244	178	198	154	190	214	311	196
Weighted base	2048	178	104	1766	86	235	174	184	151	198	270	286	180
NET: In the last year	1505 73%	134 75%	74 71%	1296 73%	65 75%	172 73%	142 82%	138 75%	111 73%	141 71%	188 69%	211 74%	129 71%
In the last month	925 45%	89 50%	42 41%	794 45%	39 45%	109 46%	94 54%	84 46%	75 49%	83 42%	100 37%	121 42%	89 49%
More than 1 month and up to 6 months ago	381 19%	24 14%	22 21%	335 19%	15 17%	46 20%	32 18%	36 20%	27 18%	39 20%	59 22%	62 22%	19 11%
More than 6 months and up to 1 year ago	198 10%	21 12%	10 10%	167 9%	11 13%	16 7%	17 10%	18 10%	9 6%	20 10%	29 11%	28 10%	20 11%
More than 1 year and up to 5 years ago	279 14%	32 18%	17 16%	230 13%	10 12%	31 13%	19 11%	20 11%	18 12%	23 12%	44 16%	36 13%	27 15%
More than 5 years ago	197 10%	10 6%	11 11%	175 10%	9 11%	27 11%	10 6%	21 11%	15 10%	25 13%	27 10%	25 9%	16 9%
I have never been prescribed any medicine on the NHS	68 3%	1 1%	2 2%	65 4%	2 2%	6 2%	2 1%	5 3%	8 5%	9 4%	11 4%	14 5%	8 5%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 3

Q.1 When was the last time you were prescribed any type of medicine on the NHS, if ever?

Base: All respondents

	Last Prescribed on NHS							Condition Have Received NHS Prescription For									Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago	I have never been prescribed any medicine on the NHS	High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	2048	1552	1018	347	187	246	179	71	460	370	244	281	223	171	87	829	445	446	225	1453
Weighted base	2048	1505	925	381	198	279	197	68	417	319	248	247	202	147	81	756	517	431	239	1453
NET: In the last year	1505	1505	925	381	198	-	-	-	388	306	206	223	192	145	78	611	272	394	159	1013
	73%	100%	100%	100%	100%	-	-	-	93%	96%	83%	90%	95%	98%	96%	81%	53%	91%	67%	70%
In the last month	925	925	925	-	-	-	-	-	304	248	149	177	142	127	65	412	76	282	101	588
	45%	62%	100%	-	-	-	-	-	73%	78%	60%	72%	70%	86%	81%	55%	15%	66%	42%	40%
More than 1 month and up to 6 months ago	381	381	-	381	-	-	-	-	77	50	39	34	34	16	12	128	118	85	36	268
	19%	25%	-	100%	-	-	-	-	19%	16%	16%	14%	17%	11%	15%	17%	23%	20%	15%	18%
More than 6 months and up to 1 year ago	198	198	-	-	198	-	-	-	7	8	18	12	16	2	*	71	78	27	22	158
	10%	13%	-	-	100%	-	-	-	2%	2%	7%	5%	8%	1%	*	9%	15%	6%	9%	11%
More than 1 year and up to 5 years ago	279	-	-	-	-	279	-	-	18	7	19	17	10	1	1	104	125	27	46	217
	14%	-	-	-	-	100%	-	-	4%	2%	8%	7%	5%	1%	1%	14%	24%	6%	19%	15%
More than 5 years ago	197	-	-	-	-	-	197	-	11	5	23	7	*	2	3	41	120	9	25	165
	10%	-	-	-	-	-	100%	-	3%	2%	9%	3%	*	1%	3%	5%	23%	2%	10%	11%
I have never been prescribed any medicine on the NHS	68	-	-	-	-	-	-	68	-	-	-	-	-	-	-	-	-	*	10	58
	3%	-	-	-	-	-	-	100%	-	-	-	-	-	-	-	-	-	*	4%	4%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 4

Q.2 Which of the following conditions, if any, have you been prescribed medication for on the NHS?

Base: All respondents who have ever been prescribed any type of medicine on the NHS

	Gender		Age							Social Grade				Employment Sector	
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	1977	1000	977	190	314	299	371	295	508	563	559	306	549	196	599
Weighted base	1980	960	1020	225	328	322	357	291	457	536	526	439	480	263	847
High Blood Pressure	417 21%	254 26%	163 16%	8 4%	11 3%	20 6%	66 19%	91 31%	221 48%	114 21%	78 15%	87 20%	138 29%	37 14%	77 9%
High Cholesterol	319 16%	209 22%	110 11%	- -	2 1%	10 3%	44 12%	80 27%	183 40%	86 16%	62 12%	67 15%	104 22%	13 5%	58 7%
Asthma	248 13%	116 12%	132 13%	36 16%	41 12%	60 19%	38 11%	32 11%	41 9%	65 12%	49 9%	51 12%	83 17%	36 14%	108 13%
Mental Health condition	247 12%	111 12%	136 13%	37 17%	49 15%	64 20%	52 14%	29 10%	16 3%	49 9%	66 13%	63 14%	69 14%	40 15%	106 13%
Arthritis	202 10%	90 9%	112 11%	9 4%	8 2%	17 5%	37 10%	51 18%	81 18%	41 8%	37 7%	43 10%	80 17%	12 5%	55 7%
Type 2 Diabetes	147 7%	95 10%	52 5%	1 *	5 1%	6 2%	20 6%	34 12%	82 18%	33 6%	32 6%	27 6%	55 11%	6 2%	23 3%
Heart Disease	81 4%	59 6%	22 2%	2 1%	- -	2 1%	8 2%	13 5%	56 12%	18 3%	24 5%	15 4%	24 5%	2 1%	14 2%
Any other condition	756 38%	345 36%	411 40%	53 23%	129 39%	120 37%	155 43%	111 38%	188 41%	212 40%	210 40%	147 34%	187 39%	75 29%	303 36%
None of the above	517 26%	233 24%	284 28%	106 47%	120 37%	99 31%	87 24%	63 22%	41 9%	138 26%	158 30%	125 28%	96 20%	97 37%	285 34%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 5

Q.2 Which of the following conditions, if any, have you been prescribed medication for on the NHS?

Base: All respondents who have ever been prescribed any type of medicine on the NHS

	Region												
	Total	Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	1977	171	109	1697	75	238	174	192	147	183	204	297	187
Weighted base	1980	177	102	1701	84	229	172	179	144	190	259	273	172
High Blood Pressure	417 21%	32 18%	23 23%	362 21%	11 14%	55 24%	34 20%	36 20%	31 22%	40 21%	44 17%	77 28%	32 19%
High Cholesterol	319 16%	23 13%	17 16%	279 16%	6 7%	36 16%	31 18%	30 17%	24 17%	35 18%	35 14%	57 21%	25 15%
Asthma	248 13%	16 9%	18 17%	214 13%	21 26%	28 12%	11 7%	21 12%	19 14%	32 17%	22 8%	37 14%	23 13%
Mental Health condition	247 12%	27 15%	11 11%	209 12%	17 20%	21 9%	34 20%	20 11%	15 10%	18 9%	29 11%	35 13%	21 12%
Arthritis	202 10%	22 12%	8 8%	172 10%	8 9%	29 12%	14 8%	21 12%	21 15%	10 6%	26 10%	30 11%	13 8%
Type 2 Diabetes	147 7%	8 5%	7 7%	132 8%	4 5%	22 9%	8 5%	16 9%	15 11%	15 8%	19 7%	20 7%	13 8%
Heart Disease	81 4%	9 5%	3 3%	68 4%	3 3%	8 3%	2 1%	2 1%	6 4%	10 5%	14 5%	17 6%	7 4%
Any other condition	756 38%	75 42%	39 38%	642 38%	32 38%	88 38%	63 37%	57 32%	58 40%	86 45%	88 34%	101 37%	69 40%
None of the above	517 26%	40 23%	28 28%	448 26%	16 19%	56 24%	51 30%	52 29%	33 23%	43 22%	96 37%	54 20%	49 29%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 6

Q.2 Which of the following conditions, if any, have you been prescribed medication for on the NHS?

Base: All respondents who have ever been prescribed any type of medicine on the NHS

	Last Prescribed on NHS							I have never been prescribed any medicine on the NHS	Condition Have Received NHS Prescription For								Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago		High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	1977	1552	1018	347	187	246	179	-	460	370	244	281	223	171	87	829	445	445	217	1390
Weighted base	1980	1505	925	381	198	279	197	-	417	319	248	247	202	147	81	756	517	431	229	1395
High Blood Pressure	417	388	304	77	7	18	11	-	417	208	49	45	96	94	50	121	-	143	24	265
	21%	26%	33%	20%	4%	6%	6%	-	100%	65%	20%	18%	48%	64%	61%	16%	-	33%	11%	19%
High Cholesterol	319	306	248	50	8	7	5	-	208	319	33	31	71	91	51	101	-	148	16	167
	16%	20%	27%	13%	4%	3%	3%	-	50%	100%	13%	12%	35%	62%	63%	13%	-	34%	7%	12%
Asthma	248	206	149	39	18	19	23	-	49	33	248	40	40	15	9	67	-	59	46	150
	13%	14%	16%	10%	9%	7%	12%	-	12%	10%	100%	16%	20%	10%	11%	9%	-	14%	20%	11%
Mental Health condition	247	223	177	34	12	17	7	-	45	31	40	247	32	13	7	76	-	94	47	125
	12%	15%	19%	9%	6%	6%	3%	-	11%	10%	16%	100%	16%	9%	9%	10%	-	22%	20%	9%
Arthritis	202	192	142	34	16	10	*	-	96	71	40	32	202	34	20	70	-	51	9	143
	10%	13%	15%	9%	8%	3%	*	-	23%	22%	16%	13%	100%	23%	24%	9%	-	12%	4%	10%
Type 2 Diabetes	147	145	127	16	2	1	2	-	94	91	15	13	34	147	22	39	-	58	11	82
	7%	10%	14%	4%	1%	*	1%	-	22%	29%	6%	5%	17%	100%	28%	5%	-	13%	5%	6%
Heart Disease	81	78	65	12	*	1	3	-	50	51	9	7	20	22	81	24	-	31	3	48
	4%	5%	7%	3%	*	*	1%	-	12%	16%	4%	3%	10%	15%	100%	3%	-	7%	1%	3%
Any other condition	756	611	412	128	71	104	41	-	121	101	67	76	70	39	24	756	-	172	80	536
	38%	41%	45%	34%	36%	37%	21%	-	29%	32%	27%	31%	35%	26%	30%	100%	-	40%	35%	38%
None of the above	517	272	76	118	78	125	120	-	-	-	-	-	-	-	-	517	-	48	62	424
	26%	18%	8%	31%	39%	45%	61%	-	-	-	-	-	-	-	-	100%	-	11%	27%	30%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 7
Q.3 When you are prescribed medication, how often do you read the instructions, if ever?
Base: All respondents

	Gender		Age							Social Grade				Employment Sector	
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	2048	1043	1005	202	341	314	379	300	512	581	582	314	571	204	631
Weighted base	2048	997	1051	238	355	336	365	295	459	551	551	446	500	274	885
NET: Always/ sometimes	1757 86%	838 84%	920 88%	195 82%	294 83%	290 86%	305 83%	266 90%	409 89%	481 87%	472 86%	386 86%	419 84%	229 84%	747 84%
Always	1155 56%	548 55%	607 58%	110 46%	189 53%	182 54%	190 52%	195 66%	289 63%	325 59%	310 56%	238 53%	282 56%	136 50%	470 53%
Sometimes	602 29%	290 29%	312 30%	85 36%	105 30%	108 32%	114 31%	71 24%	120 26%	156 28%	162 29%	148 33%	137 27%	94 34%	277 31%
Rarely	189 9%	96 10%	94 9%	27 11%	38 11%	28 8%	43 12%	18 6%	36 8%	44 8%	50 9%	46 10%	49 10%	30 11%	90 10%
Never	63 3%	38 4%	24 2%	11 5%	12 3%	10 3%	10 3%	6 2%	12 3%	17 3%	15 3%	9 2%	21 4%	9 3%	26 3%
NET: Rarely/ never	252 12%	134 13%	118 11%	38 16%	50 14%	38 11%	53 15%	24 8%	48 11%	61 11%	65 12%	56 12%	70 14%	39 14%	116 13%
Not applicable	39 2%	26 3%	13 1%	5 2%	11 3%	8 2%	7 2%	6 2%	2 *	9 2%	14 3%	5 1%	11 2%	6 2%	22 2%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 8

Q.3 When you are prescribed medication, how often do you read the instructions, if ever?

Base: All respondents

	Region												
	Total	Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	2048	174	111	1763	78	244	178	198	154	190	214	311	196
Weighted base	2048	178	104	1766	86	235	174	184	151	198	270	286	180
NET: Always/ sometimes	1757 86%	161 90%	90 86%	1506 85%	69 80%	205 87%	156 89%	151 82%	121 80%	169 85%	240 89%	237 83%	158 88%
Always	1155 56%	103 58%	62 59%	991 56%	40 46%	123 52%	106 61%	106 57%	77 51%	115 58%	157 58%	159 55%	110 61%
Sometimes	602 29%	58 33%	28 27%	516 29%	29 34%	82 35%	50 29%	45 25%	45 30%	55 27%	83 31%	78 27%	48 27%
Rarely	189 9%	12 7%	8 8%	169 10%	14 16%	24 10%	10 6%	20 11%	22 14%	22 11%	10 4%	34 12%	14 8%
Never	63 3%	3 2%	5 4%	55 3%	3 3%	4 2%	6 4%	9 5%	6 4%	3 1%	11 4%	12 4%	2 1%
NET: Rarely/ never	252 12%	15 8%	13 12%	224 13%	17 20%	28 12%	16 9%	29 16%	27 18%	25 12%	22 8%	45 16%	16 9%
Not applicable	39 2%	2 1%	1 1%	35 2%	* *	2 1%	2 1%	4 2%	3 2%	5 2%	8 3%	5 2%	6 4%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 9

Q.3 When you are prescribed medication, how often do you read the instructions, if ever?

Base: All respondents

	Last Prescribed on NHS							I have never been prescribed any medicine on the NHS	Condition Have Received NHS Prescription For								Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago		High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	2048	1552	1018	347	187	246	179	71	460	370	244	281	223	171	87	829	445	446	225	1453
Weighted base	2048	1505	925	381	198	279	197	68	417	319	248	247	202	147	81	756	517	431	239	1453
NET: Always/ sometimes	1757 86%	1302 87%	790 85%	344 90%	168 85%	251 90%	179 91%	26 38%	351 84%	269 84%	207 84%	201 81%	184 91%	117 79%	68 84%	686 91%	467 90%	374 87%	182 76%	1265 87%
Always	1155 56%	834 55%	505 55%	214 56%	115 58%	167 60%	140 71%	15 22%	223 54%	184 58%	124 50%	105 42%	114 56%	80 54%	45 56%	469 62%	325 63%	220 51%	114 48%	858 59%
Sometimes	602 29%	468 31%	285 31%	130 34%	53 27%	84 30%	39 20%	11 16%	127 30%	85 27%	83 34%	96 39%	70 35%	37 25%	23 28%	217 29%	142 27%	154 36%	68 29%	407 28%
Rarely	189 9%	149 10%	103 11%	27 7%	19 10%	20 7%	12 6%	9 13%	49 12%	37 12%	31 13%	36 15%	11 6%	26 17%	8 9%	58 8%	25 5%	47 11%	45 19%	108 7%
Never	63 3%	49 3%	30 3%	9 2%	9 5%	3 1%	3 2%	8 11%	17 4%	13 4%	9 4%	9 3%	7 3%	5 3%	5 6%	11 2%	15 3%	10 2%	12 5%	41 3%
NET: Rarely/ never	252 12%	198 13%	134 14%	36 10%	28 14%	23 8%	15 8%	16 24%	66 16%	50 16%	41 16%	45 18%	18 9%	31 21%	13 16%	70 9%	40 8%	57 13%	57 24%	149 10%
Not applicable	39 2%	4 *	2 *	1 *	1 1%	5 2%	3 1%	26 38%	* *	- -	- -	1 *	- -	- -	1 1%	1 *	10 2%	- -	- -	39 3%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 10

Q.4 Thinking about prescription medication, which of the following actions, if any, have you taken in the past?

Base: All respondents

	Gender			Age						Social Grade				Employment Sector	
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	2048	1043	1005	202	341	314	379	300	512	581	582	314	571	204	631
Weighted base	2048	997	1051	238	355	336	365	295	459	551	551	446	500	274	885
Disposed of medication that you did not use	625 31%	318 32%	307 29%	60 25%	116 33%	114 34%	104 28%	89 30%	143 31%	194 35%	150 27%	146 33%	136 27%	105 38%	259 29%
Kept old drugs that you had not taken	554 27%	222 22%	332 32%	108 45%	106 30%	95 28%	95 26%	70 24%	80 17%	188 34%	154 28%	112 25%	101 20%	100 36%	250 28%
Returned medication that you did not use to your doctor, healthcare professional or pharmacy	520 25%	236 24%	284 27%	15 6%	38 11%	56 17%	104 29%	97 33%	209 46%	127 23%	137 25%	109 24%	147 29%	58 21%	126 14%
Missed a round of medication that you are meant to take every day, such as statins for high cholesterol	431 21%	224 22%	207 20%	49 20%	55 15%	48 14%	95 26%	65 22%	120 26%	123 22%	117 21%	89 20%	102 20%	61 22%	170 19%
Did not finish a course of medication, such as antibiotics	239 12%	105 11%	134 13%	41 17%	83 23%	49 15%	35 10%	11 4%	21 5%	72 13%	66 12%	52 12%	48 10%	47 17%	129 15%
Given medication that you had not taken to someone you know	120 6%	46 5%	74 7%	22 9%	30 8%	21 6%	19 5%	18 6%	11 2%	34 6%	17 3%	37 8%	31 6%	27 10%	58 7%
Informed your doctor or healthcare professional that you did not finish a course of medication	119 6%	64 6%	55 5%	21 9%	31 9%	20 6%	14 4%	14 5%	19 4%	47 9%	28 5%	22 5%	21 4%	17 6%	54 6%
None of these	632 31%	326 33%	306 29%	64 27%	114 32%	127 38%	105 29%	96 32%	126 27%	143 26%	188 34%	140 31%	162 32%	74 27%	297 34%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 11

Q.4 Thinking about prescription medication, which of the following actions, if any, have you taken in the past?

Base: All respondents

	Region												
	Total	Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	2048	174	111	1763	78	244	178	198	154	190	214	311	196
Weighted base	2048	178	104	1766	86	235	174	184	151	198	270	286	180
Disposed of medication that you did not use	625 31%	52 29%	35 33%	539 31%	22 25%	71 30%	51 29%	60 32%	43 28%	70 35%	87 32%	87 30%	49 27%
Kept old drugs that you had not taken	554 27%	38 22%	25 24%	491 28%	33 38%	58 24%	57 33%	38 20%	34 22%	60 30%	76 28%	85 30%	51 28%
Returned medication that you did not use to your doctor, healthcare professional or pharmacy	520 25%	62 35%	24 23%	433 25%	11 13%	45 19%	45 26%	47 26%	46 30%	61 31%	51 19%	77 27%	50 28%
Missed a round of medication that you are meant to take every day, such as statins for high cholesterol	431 21%	34 19%	24 23%	373 21%	19 22%	56 24%	44 25%	26 14%	34 23%	46 23%	46 17%	70 24%	33 18%
Did not finish a course of medication, such as antibiotics	239 12%	21 12%	11 10%	208 12%	10 12%	19 8%	22 13%	12 6%	21 14%	25 12%	43 16%	42 15%	14 8%
Given medication that you had not taken to someone you know	120 6%	17 10%	13 13%	89 5%	6 6%	11 5%	10 6%	8 4%	9 6%	6 3%	14 5%	18 6%	7 4%
Informed your doctor or healthcare professional that you did not finish a course of medication	119 6%	13 8%	5 4%	101 6%	4 4%	9 4%	11 6%	6 3%	9 6%	16 8%	26 10%	12 4%	8 4%
None of these	632 31%	52 29%	34 33%	547 31%	27 31%	81 34%	41 24%	62 34%	45 30%	52 26%	88 33%	87 30%	63 35%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 12

Q.4 Thinking about prescription medication, which of the following actions, if any, have you taken in the past?

Base: All respondents

	Last Prescribed on NHS							I have never been prescribed any medicine on the NHS	Condition Have Received NHS Prescription For								Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago		High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	2048	1552	1018	347	187	246	179	71	460	370	244	281	223	171	87	829	445	446	225	1453
Weighted base	2048	1505	925	381	198	279	197	68	417	319	248	247	202	147	81	756	517	431	239	1453
Disposed of medication that you did not use	625 31%	476 32%	297 32%	114 30%	66 33%	92 33%	50 25%	8 11%	130 31%	119 37%	90 36%	94 38%	64 31%	52 35%	25 31%	243 32%	152 29%	188 44%	109 45%	372 26%
Kept old drugs that you had not taken	554 27%	439 29%	261 28%	123 32%	54 27%	79 28%	33 17%	4 5%	84 20%	63 20%	95 38%	115 47%	53 26%	29 20%	12 14%	227 30%	127 25%	182 42%	110 46%	307 21%
Returned medication that you did not use to your doctor, healthcare professional or pharmacy	520 25%	447 30%	348 38%	66 17%	33 17%	48 17%	22 11%	3 4%	168 40%	147 46%	59 24%	73 30%	89 44%	73 50%	46 57%	257 34%	53 10%	149 35%	43 18%	347 24%
Missed a round of medication that you are meant to take every day, such as statins for high cholesterol	431 21%	394 26%	282 31%	85 22%	27 14%	27 10%	9 5%	* *	143 34%	148 46%	59 24%	94 38%	51 25%	58 39%	31 38%	172 23%	48 9%	431 100%	75 32%	- -
Did not finish a course of medication, such as antibiotics	239 12%	159 11%	101 11%	36 9%	22 11%	46 16%	25 13%	10 14%	24 6%	16 5%	46 19%	47 19%	9 5%	11 7%	3 3%	80 11%	62 12%	75 17%	239 100%	- -
Given medication that you had not taken to someone you know	120 6%	106 7%	60 6%	36 9%	11 5%	9 3%	3 2%	1 2%	30 7%	12 4%	19 8%	39 16%	19 9%	7 5%	3 4%	47 6%	19 4%	47 11%	45 19%	54 4%
Informed your doctor or healthcare professional that you did not finish a course of medication	119 6%	109 7%	72 8%	22 6%	14 7%	6 2%	3 2%	1 2%	29 7%	17 5%	22 9%	32 13%	20 10%	9 6%	2 2%	40 5%	20 4%	38 9%	36 15%	64 4%
None of these	632 31%	386 26%	210 23%	114 30%	62 31%	101 36%	98 50%	47 69%	107 26%	65 20%	57 23%	35 14%	38 19%	30 20%	21 26%	191 25%	211 41%	- -	- -	632 43%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 13

Q.5 What are the main reasons that have stopped you from taking all of your medication as prescribed in the past?

Base: All respondents who have failed to finish or missed a round of prescription medication

	Gender		Age						Social Grade				Employment Sector		
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	595	323	272	65	106	90	109	78	147	179	169	78	169	67	184
Weighted base	595	297	298	79	116	87	110	73	131	176	164	120	135	95	257
I forgot to take the medication	385 65%	185 62%	200 67%	53 68%	67 58%	50 57%	69 63%	54 73%	93 71%	117 66%	110 67%	66 55%	93 69%	61 64%	158 61%
The medication had side-effects that made me feel unwell	149 25%	62 21%	87 29%	20 25%	20 17%	26 30%	37 34%	13 18%	34 26%	42 24%	42 26%	35 29%	30 22%	32 34%	52 20%
I didn't feel ill and therefore didn't think I needed the medication	118 20%	61 21%	57 19%	22 27%	38 33%	25 29%	14 13%	8 11%	10 8%	40 23%	33 20%	27 23%	17 12%	18 19%	75 29%
I worried about the side-effects of taking medication everyday	56 9%	28 9%	29 10%	3 3%	12 11%	9 10%	7 6%	15 21%	11 8%	15 8%	19 12%	15 13%	7 5%	7 7%	28 11%
I wanted to be able to have an alcoholic drink	42 7%	22 8%	20 7%	8 10%	18 15%	6 7%	8 7%	1 2%	1 1%	12 7%	15 9%	8 7%	7 5%	15 16%	21 8%
The cost of the prescription was too high	40 7%	14 5%	26 9%	11 14%	10 8%	7 8%	10 9%	2 2%	- -	19 11%	10 6%	5 4%	6 4%	10 11%	20 8%
I had multiple prescriptions and lost track of what needed to be taken when	31 5%	19 6%	12 4%	- -	6 5%	8 9%	6 6%	2 2%	9 7%	8 5%	5 3%	4 3%	13 10%	6 7%	7 3%
The course of medication that I had to take was too long	21 4%	11 4%	10 3%	3 4%	7 6%	5 6%	4 4%	1 1%	1 1%	6 3%	5 3%	5 4%	5 4%	5 6%	12 5%
The instructions on how and when to take it were too complicated	6 1%	4 1%	2 1%	- -	2 2%	2 3%	- -	1 1%	1 1%	3 2%	2 1%	- -	1 1%	2 2%	2 1%
None of the above	32 5%	22 7%	10 3%	9 12%	1 1%	5 5%	1 1%	5 6%	12 9%	6 3%	9 5%	9 8%	9 6%	1 1%	13 5%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 14

Q.5 What are the main reasons that have stopped you from taking all of your medication as prescribed in the past?

Base: All respondents who have failed to finish or missed a round of prescription medication

	Total	Region											
		Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	595	44	29	522	25	83	55	40	49	58	63	96	53
Weighted base	595	48	30	517	29	67	58	35	53	63	76	95	42
I forgot to take the medication	385 65%	31 64%	22 74%	332 64%	17 59%	49 72%	35 61%	22 64%	31 59%	44 70%	45 60%	59 62%	30 71%
The medication had side-effects that made me feel unwell	149 25%	7 15%	7 22%	135 26%	6 21%	15 22%	22 39%	8 24%	14 28%	16 26%	18 24%	21 22%	13 31%
I didn't feel ill and therefore didn't think I needed the medication	118 20%	14 30%	3 11%	100 19%	8 28%	9 13%	1 2%	7 20%	8 14%	17 27%	24 32%	19 20%	7 17%
I worried about the side-effects of taking medication everyday	56 9%	2 4%	2 5%	53 10%	1 3%	9 14%	6 11%	5 14%	2 4%	5 8%	11 14%	11 12%	3 6%
I wanted to be able to have an alcoholic drink	42 7%	4 8%	4 14%	35 7%	1 5%	6 9%	8 14%	1 3%	*	1 1%	8 11%	7 7%	2 4%
The cost of the prescription was too high	40 7%	-	*	39 8%	4 12%	5 7%	5 9%	5 13%	1 2%	7 11%	*	10 10%	3 8%
I had multiple prescriptions and lost track of what needed to be taken when	31 5%	3 5%	4 14%	24 5%	1 4%	4 6%	2 4%	3 10%	4 7%	1 1%	2 3%	4 5%	1 3%
The course of medication that I had to take was too long	21 4%	1 3%	2 8%	17 3%	-	3 4%	5 9%	-	4 7%	-	3 4%	1 1%	2 4%
The instructions on how and when to take it were too complicated	6 1%	-	-	6 1%	-	1 2%	-	*	-	2 4%	1 1%	-	1 3%
None of the above	32 5%	1 2%	2 5%	30 6%	-	2 3%	2 4%	3 10%	6 11%	4 6%	3 3%	7 7%	2 5%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 15

Q.5 What are the main reasons that have stopped you from taking all of your medication as prescribed in the past?

Base: All respondents who have failed to finish or missed a round of prescription medication

	Last Prescribed on NHS							Condition Have Received NHS Prescription For									Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago	I have never been prescribed any medicine on the NHS	High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	595	505	363	100	42	54	28	8	167	174	90	135	71	75	35	249	79	446	225	-
Weighted base	595	492	338	113	40	62	32	10	152	152	97	122	59	65	33	220	93	431	239	-
I forgot to take the medication	385 65%	328 67%	240 71%	66 58%	22 55%	40 65%	16 49%	1 15%	115 75%	107 70%	62 64%	81 66%	45 77%	48 74%	29 88%	147 67%	53 57%	313 73%	122 51%	-
The medication had side-effects that made me feel unwell	149 25%	132 27%	103 30%	24 21%	5 13%	9 15%	5 16%	2 25%	33 22%	41 27%	30 31%	42 35%	15 25%	12 18%	5 16%	55 25%	19 21%	117 27%	65 27%	-
I didn't feel ill and therefore didn't think I needed the medication	118 20%	81 17%	43 13%	22 20%	16 40%	16 26%	18 57%	2 18%	10 7%	10 6%	17 17%	25 20%	6 10%	7 11%	3 9%	35 16%	30 33%	51 12%	90 38%	-
I worried about the side-effects of taking medication everyday	56 9%	51 10%	27 8%	18 16%	6 14%	4 6%	2 5%	* 3%	14 9%	18 12%	10 10%	14 11%	6 10%	2 4%	7 20%	21 9%	10 10%	45 10%	19 8%	-
I wanted to be able to have an alcoholic drink	42 7%	37 7%	18 5%	11 10%	8 20%	3 4%	2 5%	1 12%	4 2%	6 4%	6 6%	6 5%	- -	2 3%	- -	10 4%	13 14%	24 5%	26 11%	-
The cost of the prescription was too high	40 7%	24 5%	13 4%	8 7%	4 9%	8 13%	7 22%	- -	5 3%	7 4%	11 11%	11 9%	- -	- -	- -	9 4%	7 7%	25 6%	22 9%	-
I had multiple prescriptions and lost track of what needed to be taken when	31 5%	31 6%	26 8%	4 3%	1 2%	- -	- -	- -	8 5%	9 6%	7 7%	9 8%	7 11%	5 8%	1 4%	12 5%	3 3%	27 6%	8 3%	-
The course of medication that I had to take was too long	21 4%	16 3%	7 2%	8 7%	1 3%	4 7%	- -	1 10%	3 2%	1 1%	4 4%	5 4%	1 1%	1 2%	- -	4 2%	5 6%	10 2%	18 8%	-
The instructions on how and when to take it were too complicated	6 1%	6 1%	1 *	1 1%	3 8%	- -	- -	- -	* *	1 1%	- -	2 2%	* 1%	1 1%	- -	1 *	1 2%	5 1%	1 *	-
None of the above	32 5%	28 6%	15 4%	10 9%	4 9%	1 2%	- -	2 23%	10 7%	14 9%	3 3%	3 3%	2 3%	6 9%	1 2%	9 4%	5 5%	28 6%	8 3%	-

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 16

Q.6 If you had medication left over from a prescription that you had not finished, what would you be most likely to do with it?

Base: All respondents

	Gender		Age						Social Grade				Employment Sector		
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	2048	1043	1005	202	341	314	379	300	512	581	582	314	571	204	631
Weighted base	2048	997	1051	238	355	336	365	295	459	551	551	446	500	274	885
Take it to my local pharmacy	693 34%	332 33%	361 34%	34 14%	55 15%	83 25%	126 35%	139 47%	256 56%	154 28%	198 36%	154 35%	186 37%	75 27%	199 23%
Keep it in case I need it later	507 25%	202 20%	305 29%	92 39%	119 33%	78 23%	87 24%	67 23%	64 14%	157 29%	130 24%	108 24%	112 22%	72 26%	266 30%
Throw it away	494 24%	270 27%	224 21%	69 29%	121 34%	119 35%	85 23%	43 15%	56 12%	156 28%	137 25%	101 23%	101 20%	82 30%	262 30%
Flush it down the toilet	133 6%	78 8%	55 5%	9 4%	15 4%	14 4%	32 9%	23 8%	40 9%	36 7%	20 4%	38 9%	39 8%	17 6%	55 6%
Take it to my GP or other healthcare professional	120 6%	72 7%	48 5%	10 4%	16 5%	19 6%	26 7%	15 5%	33 7%	31 6%	29 5%	26 6%	34 7%	14 5%	55 6%
Give it to a friend or family member that might need it	21 1%	7 1%	14 1%	4 2%	7 2%	8 2%	1 *	1 *	- -	4 1%	4 1%	10 2%	3 1%	4 2%	13 2%
Don't know	79 4%	37 4%	42 4%	19 8%	22 6%	14 4%	8 2%	6 2%	10 2%	13 2%	32 6%	10 2%	25 5%	10 4%	34 4%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 17

Q.6 If you had medication left over from a prescription that you had not finished, what would you be most likely to do with it?

Base: All respondents

	Region												
	Total	Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	2048	174	111	1763	78	244	178	198	154	190	214	311	196
Weighted base	2048	178	104	1766	86	235	174	184	151	198	270	286	180
Take it to my local pharmacy	693 34%	73 41%	29 28%	591 33%	22 25%	75 32%	62 36%	67 36%	60 40%	71 36%	54 20%	113 39%	67 37%
Keep it in case I need it later	507 25%	39 22%	22 21%	447 25%	34 39%	46 20%	47 27%	48 26%	35 23%	47 24%	76 28%	69 24%	45 25%
Throw it away	494 24%	37 21%	25 24%	433 25%	21 24%	55 24%	41 23%	47 26%	31 20%	42 21%	92 34%	64 22%	40 22%
Flush it down the toilet	133 6%	12 7%	13 13%	107 6%	3 3%	27 12%	10 6%	9 5%	9 6%	17 8%	10 4%	11 4%	12 6%
Take it to my GP or other healthcare professional	120 6%	10 6%	7 7%	103 6%	2 3%	17 7%	9 5%	8 4%	10 7%	18 9%	16 6%	15 5%	7 4%
Give it to a friend or family member that might need it	21 1%	6 3%	2 2%	13 1%	3 3%	1 *	- -	1 *	1 1%	- -	5 2%	4 1%	- -
Don't know	79 4%	2 1%	6 6%	71 4%	2 2%	13 6%	4 2%	5 3%	6 4%	4 2%	18 7%	11 4%	9 5%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 18

Q.6 If you had medication left over from a prescription that you had not finished, what would you be most likely to do with it?

Base: All respondents

	Last Prescribed on NHS							I have never been prescribed any medicine on the NHS	Condition Have Received NHS Prescription For								Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago		High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	2048	1552	1018	347	187	246	179	71	460	370	244	281	223	171	87	829	445	446	225	1453
Weighted base	2048	1505	925	381	198	279	197	68	417	319	248	247	202	147	81	756	517	431	239	1453
Take it to my local pharmacy	693 34%	558 37%	404 44%	101 27%	53 27%	67 24%	60 31%	8 11%	191 46%	154 48%	82 33%	80 32%	97 48%	65 44%	42 52%	303 40%	115 22%	141 33%	43 18%	520 36%
Keep it in case I need it later	507 25%	363 24%	211 23%	99 26%	53 27%	81 29%	49 25%	14 21%	63 15%	42 13%	67 27%	77 31%	34 17%	19 13%	7 9%	191 25%	147 28%	123 28%	94 39%	329 23%
Throw it away	494 24%	315 21%	152 16%	102 27%	61 31%	102 37%	62 32%	14 21%	76 18%	53 17%	70 28%	58 24%	27 13%	23 16%	13 16%	151 20%	166 32%	92 21%	80 34%	340 23%
Flush it down the toilet	133 6%	101 7%	48 5%	41 11%	12 6%	12 4%	17 9%	2 2%	41 10%	32 10%	12 5%	8 3%	20 10%	14 10%	4 5%	45 6%	42 8%	35 8%	8 4%	92 6%
Take it to my GP or other healthcare professional	120 6%	106 7%	75 8%	21 5%	10 5%	6 2%	3 1%	6 8%	34 8%	32 10%	9 4%	11 4%	15 7%	20 14%	12 15%	52 7%	20 4%	31 7%	4 2%	86 6%
Give it to a friend or family member that might need it	21 1%	18 1%	13 1%	3 1%	2 1%	1 *	1 1%	* *	4 1%	* *	1 *	7 3%	2 1%	- -	- -	1 *	6 1%	5 1%	5 2%	15 1%
Don't know	79 4%	43 3%	21 2%	14 4%	7 4%	9 3%	4 2%	24 35%	9 2%	4 1%	6 2%	6 3%	7 4%	6 4%	2 2%	14 2%	22 4%	5 1%	5 2%	70 5%

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 19

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

Summary**Base: All respondents**

	Aspects		
	Having personalised packaging from your pharmacist that shows when each dose should be taken	Being sent reminders by text or by phone	The cost of the medicine to the NHS being printed on the packaging
Unweighted base	2048	2048	2048
Weighted base	2048	2048	2048
NET: More likely	1078 53%	835 41%	743 36%
Much more likely (+2)	453 22%	332 16%	284 14%
Slightly more likely (+1)	625 31%	503 25%	459 22%
Neither more likely nor less likely (0)	812 40%	893 44%	1013 49%
Slightly less likely (-1)	27 1%	65 3%	54 3%
Much less likely (-2)	15 1%	67 3%	45 2%
NET: Less likely	42 2%	132 6%	99 5%
Don't know/ not applicable	116 6%	189 9%	192 9%
Mean	0.76	0.52	0.48
Standard deviation	0.85	0.95	0.87
Standard error	0.02	0.02	0.02

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 20

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

The cost of the medicine to the NHS being printed on the packaging

Base: All respondents

	Gender			Age						Social Grade				Employment Sector	
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	2048	1043	1005	202	341	314	379	300	512	581	582	314	571	204	631
Weighted base	2048	997	1051	238	355	336	365	295	459	551	551	446	500	274	885
NET: More likely	743 36%	342 34%	401 38%	99 42%	139 39%	127 38%	141 39%	85 29%	152 33%	226 41%	183 33%	177 40%	158 32%	124 45%	336 38%
Much more likely	(+2) 284 14%	124 12%	161 15%	34 14%	55 16%	51 15%	44 12%	30 10%	69 15%	82 15%	76 14%	67 15%	60 12%	56 20%	114 13%
Slightly more likely	(+1) 459 22%	219 22%	240 23%	65 27%	84 24%	76 22%	96 26%	55 19%	83 18%	144 26%	107 19%	110 25%	98 20%	68 25%	222 25%
Neither more likely nor less likely	(0) 1013 49%	504 51%	509 48%	87 36%	147 42%	170 50%	188 51%	172 58%	250 54%	267 48%	297 54%	191 43%	259 52%	110 40%	423 48%
Slightly less likely	(-1) 54 3%	36 4%	18 2%	15 6%	9 2%	8 2%	6 2%	10 4%	7 2%	11 2%	9 2%	16 4%	18 4%	7 3%	31 3%
Much less likely	(-2) 45 2%	24 2%	21 2%	7 3%	17 5%	5 2%	4 1%	5 2%	7 1%	11 2%	13 2%	5 1%	16 3%	5 2%	24 3%
NET: Less likely	99 5%	60 6%	39 4%	22 9%	25 7%	13 4%	10 3%	16 5%	14 3%	22 4%	22 4%	21 5%	34 7%	12 4%	55 6%
Don't know/ not applicable	192 9%	91 9%	101 10%	30 13%	43 12%	26 8%	27 7%	22 7%	44 10%	36 7%	49 9%	59 13%	48 10%	28 10%	71 8%
Mean	0.48	0.42	0.53	0.50	0.49	0.52	0.50	0.35	0.48	0.53	0.44	0.56	0.37	0.66	0.46
Standard deviation	0.87	0.87	0.87	0.97	0.99	0.86	0.79	0.80	0.84	0.86	0.86	0.87	0.89	0.93	0.88
Standard error	0.02	0.03	0.03	0.07	0.06	0.05	0.04	0.05	0.04	0.04	0.04	0.05	0.04	0.07	0.04

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 21

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

The cost of the medicine to the NHS being printed on the packaging

Base: All respondents

	Total	Region											
		Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	2048	174	111	1763	78	244	178	198	154	190	214	311	196
Weighted base	2048	178	104	1766	86	235	174	184	151	198	270	286	180
NET: More likely	743 36%	79 44%	45 43%	620 35%	24 28%	76 32%	65 37%	72 39%	50 33%	69 35%	103 38%	90 31%	70 39%
Much more likely (+2)	284 14%	28 16%	19 19%	237 13%	12 14%	26 11%	26 15%	34 18%	26 17%	16 8%	39 14%	38 13%	20 11%
Slightly more likely (+1)	459 22%	51 29%	25 24%	383 22%	12 14%	51 21%	39 22%	38 21%	24 16%	53 27%	64 24%	52 18%	50 28%
Neither more likely nor less likely (0)	1013 49%	75 42%	51 48%	887 50%	55 64%	117 50%	88 50%	78 42%	84 56%	101 51%	117 43%	161 56%	87 48%
Slightly less likely (-1)	54 3%	3 2%	3 3%	48 3%	2 3%	14 6%	2 1%	3 1%	5 3%	5 3%	10 4%	3 1%	3 2%
Much less likely (-2)	45 2%	4 2%	1 1%	41 2%	- -	7 3%	3 2%	6 3%	- -	2 1%	12 4%	4 1%	7 4%
NET: Less likely	99 5%	7 4%	4 4%	88 5%	2 3%	21 9%	6 3%	9 5%	5 3%	7 4%	22 8%	7 2%	10 6%
Don't know/ not applicable	192 9%	17 10%	5 5%	170 10%	4 5%	21 9%	15 9%	26 14%	13 8%	21 11%	27 10%	29 10%	13 7%
Mean	0.48	0.59	0.60	0.46	0.42	0.35	0.52	0.58	0.51	0.43	0.44	0.45	0.44
Standard deviation	0.87	0.88	0.86	0.87	0.78	0.89	0.86	0.97	0.84	0.74	0.98	0.81	0.88
Standard error	0.02	0.07	0.08	0.02	0.09	0.06	0.07	0.07	0.07	0.06	0.07	0.05	0.07

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 22

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

The cost of the medicine to the NHS being printed on the packaging

Base: All respondents

	Last Prescribed on NHS								Condition Have Received NHS Prescription For								Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago	I have never been prescribed any medicine on the NHS	High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	2048	1552	1018	347	187	246	179	71	460	370	244	281	223	171	87	829	445	446	225	1453
Weighted base	2048	1505	925	381	198	279	197	68	417	319	248	247	202	147	81	756	517	431	239	1453
NET: More likely	743	547	298	153	96	105	77	14	123	94	89	83	63	50	22	248	224	142	91	536
	36%	36%	32%	40%	49%	38%	39%	21%	29%	29%	36%	34%	31%	34%	28%	33%	43%	33%	38%	37%
Much more likely (+2)	284	222	133	52	37	37	22	4	57	39	37	38	32	21	5	84	87	50	31	209
	14%	15%	14%	14%	19%	13%	11%	6%	14%	12%	15%	15%	16%	14%	6%	11%	17%	12%	13%	14%
Slightly more likely (+1)	459	325	164	101	59	68	55	10	66	54	51	46	32	28	17	164	137	92	59	327
	22%	22%	18%	27%	30%	24%	28%	15%	16%	17%	21%	18%	16%	19%	21%	22%	27%	21%	25%	23%
Neither more likely nor less likely (0)	1013	746	494	171	81	140	92	35	233	185	122	129	113	74	38	404	219	231	113	705
	49%	50%	53%	45%	41%	50%	47%	51%	56%	58%	49%	52%	56%	50%	47%	53%	42%	54%	47%	49%
Slightly less likely (-1)	54	39	16	20	4	5	6	4	15	4	16	7	9	4	6	14	10	7	7	45
	3%	3%	2%	5%	2%	2%	3%	6%	4%	1%	6%	3%	4%	3%	7%	2%	2%	2%	3%	3%
Much less likely (-2)	45	39	31	4	4	2	3	1	16	4	6	8	4	4	2	17	9	14	10	25
	2%	3%	3%	1%	2%	1%	2%	2%	4%	1%	2%	3%	2%	2%	3%	2%	2%	3%	4%	2%
NET: Less likely	99	78	46	24	8	8	9	5	31	9	22	15	13	7	8	31	19	21	17	70
	5%	5%	5%	6%	4%	3%	4%	8%	7%	3%	9%	6%	6%	5%	10%	4%	4%	5%	7%	5%
Don't know/ not applicable	192	134	87	33	14	26	19	14	31	31	15	20	13	17	13	72	55	37	19	142
	9%	9%	9%	9%	7%	9%	9%	20%	7%	10%	6%	8%	6%	11%	16%	10%	11%	9%	8%	10%
Mean	0.48	0.48	0.42	0.51	0.66	0.52	0.49	0.21	0.35	0.42	0.42	0.43	0.41	0.46	0.25	0.42	0.61	0.40	0.43	0.50
Standard deviation	0.87	0.89	0.90	0.86	0.90	0.80	0.82	0.77	0.92	0.80	0.93	0.92	0.90	0.90	0.85	0.83	0.88	0.86	0.93	0.87
Standard error	0.02	0.02	0.03	0.05	0.07	0.05	0.06	0.10	0.04	0.04	0.06	0.06	0.06	0.07	0.10	0.03	0.04	0.04	0.06	0.02

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 23

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

Being sent reminders by text or by phone

Base: All respondents

	Gender			Age						Social Grade				Employment Sector	
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	2048	1043	1005	202	341	314	379	300	512	581	582	314	571	204	631
Weighted base	2048	997	1051	238	355	336	365	295	459	551	551	446	500	274	885
NET: More likely	835 41%	400 40%	435 41%	147 62%	166 47%	138 41%	169 46%	89 30%	127 28%	231 42%	223 41%	196 44%	184 37%	145 53%	412 47%
Much more likely	(+2) 332 16%	153 15%	179 17%	65 27%	76 22%	48 14%	57 16%	40 14%	46 10%	85 15%	99 18%	86 19%	62 12%	59 21%	158 18%
Slightly more likely	(+1) 503 25%	247 25%	256 24%	82 34%	89 25%	90 27%	112 31%	49 16%	81 18%	147 27%	125 23%	109 25%	122 24%	87 32%	254 29%
Neither more likely nor less likely	(0) 893 44%	439 44%	454 43%	60 25%	128 36%	156 46%	142 39%	155 53%	251 55%	247 45%	244 44%	172 39%	229 46%	84 30%	350 40%
Slightly less likely	(-1) 65 3%	32 3%	33 3%	4 2%	7 2%	14 4%	10 3%	13 4%	17 4%	17 3%	13 2%	20 4%	15 3%	16 6%	25 3%
Much less likely	(-2) 67 3%	37 4%	30 3%	6 3%	13 4%	12 4%	10 3%	9 3%	18 4%	16 3%	22 4%	9 2%	21 4%	11 4%	21 2%
NET: Less likely	132 6%	69 7%	63 6%	10 4%	20 6%	26 8%	19 5%	22 8%	34 7%	33 6%	35 6%	29 6%	35 7%	27 10%	46 5%
Don't know/ not applicable	189 9%	90 9%	99 9%	20 9%	41 12%	17 5%	35 10%	29 10%	46 10%	39 7%	49 9%	50 11%	51 10%	19 7%	76 9%
Mean	0.52	0.49	0.55	0.90	0.67	0.46	0.60	0.37	0.29	0.52	0.53	0.62	0.42	0.65	0.62
Standard deviation	0.95	0.95	0.94	0.95	1.00	0.93	0.91	0.92	0.88	0.91	0.98	0.95	0.93	1.03	0.92
Standard error	0.02	0.03	0.03	0.07	0.06	0.05	0.05	0.06	0.04	0.04	0.04	0.06	0.04	0.08	0.04

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 24

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

Being sent reminders by text or by phone

Base: All respondents

	Region												
	Total	Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	2048	174	111	1763	78	244	178	198	154	190	214	311	196
Weighted base	2048	178	104	1766	86	235	174	184	151	198	270	286	180
NET: More likely	835 41%	81 45%	49 47%	705 40%	36 42%	107 45%	79 45%	81 44%	53 35%	62 31%	121 45%	99 35%	68 38%
Much more likely (+2)	332 16%	40 22%	21 20%	272 15%	16 18%	41 17%	31 18%	40 22%	20 13%	24 12%	44 16%	39 14%	17 10%
Slightly more likely (+1)	503 25%	41 23%	28 27%	434 25%	20 23%	66 28%	48 28%	40 22%	33 22%	38 19%	77 29%	60 21%	50 28%
Neither more likely nor less likely (0)	893 44%	78 44%	43 42%	771 44%	44 52%	88 37%	72 42%	68 37%	66 43%	102 51%	92 34%	149 52%	90 50%
Slightly less likely (-1)	65 3%	3 2%	2 2%	60 3%	3 4%	7 3%	2 1%	7 4%	15 10%	9 4%	10 4%	4 1%	3 2%
Much less likely (-2)	67 3%	6 4%	3 3%	58 3%	- -	10 4%	3 2%	9 5%	4 3%	4 2%	13 5%	9 3%	5 3%
NET: Less likely	132 6%	9 5%	5 4%	118 7%	3 4%	17 7%	5 3%	17 9%	19 13%	13 6%	23 8%	13 4%	9 5%
Don't know/ not applicable	189 9%	10 6%	8 7%	171 10%	3 3%	23 10%	18 10%	19 10%	13 9%	22 11%	34 13%	26 9%	14 8%
Mean	0.52	0.62	0.64	0.50	0.58	0.57	0.65	0.58	0.35	0.40	0.55	0.44	0.43
Standard deviation	0.95	0.98	0.94	0.94	0.84	1.00	0.87	1.08	0.96	0.87	1.02	0.89	0.83
Standard error	0.02	0.08	0.09	0.02	0.10	0.07	0.07	0.08	0.08	0.07	0.07	0.05	0.06

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 25

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

Being sent reminders by text or by phone

Base: All respondents

	Last Prescribed on NHS								Condition Have Received NHS Prescription For								Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago	I have never been prescribed any medicine on the NHS	High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	2048	1552	1018	347	187	246	179	71	460	370	244	281	223	171	87	829	445	446	225	1453
Weighted base	2048	1505	925	381	198	279	197	68	417	319	248	247	202	147	81	756	517	431	239	1453
NET: More likely	835	626	360	166	99	104	85	19	130	105	120	108	86	56	20	264	248	199	136	542
	41%	42%	39%	44%	50%	37%	43%	28%	31%	33%	48%	44%	42%	38%	25%	35%	48%	46%	57%	37%
Much more likely (+2)	332	263	152	73	38	36	27	6	48	42	44	46	28	27	5	112	102	79	50	218
	16%	17%	16%	19%	19%	13%	14%	9%	11%	13%	18%	19%	14%	18%	6%	15%	20%	18%	21%	15%
Slightly more likely (+1)	503	363	209	93	61	68	58	14	83	62	76	62	57	29	15	153	146	120	85	324
	25%	24%	23%	24%	31%	24%	30%	20%	20%	20%	31%	25%	28%	20%	18%	20%	28%	28%	36%	22%
Neither more likely nor less likely (0)	893	653	414	162	77	137	75	28	209	164	94	114	90	58	46	377	185	183	83	654
	44%	43%	45%	43%	39%	49%	38%	41%	50%	51%	38%	46%	44%	40%	56%	50%	36%	42%	35%	45%
Slightly less likely (-1)	65	39	23	12	5	6	12	7	18	13	11	5	5	8	5	12	14	11	5	49
	3%	3%	2%	3%	3%	2%	6%	10%	4%	4%	5%	2%	3%	5%	6%	2%	3%	3%	2%	3%
Much less likely (-2)	67	54	41	10	3	6	6	2	30	10	9	6	6	6	2	31	11	10	8	50
	3%	4%	4%	3%	1%	2%	3%	2%	7%	3%	3%	2%	3%	4%	3%	4%	2%	2%	3%	3%
NET: Less likely	132	94	64	22	8	12	18	9	47	23	20	12	11	14	7	42	25	21	13	99
	6%	6%	7%	6%	4%	4%	9%	13%	11%	7%	8%	5%	6%	10%	9%	6%	5%	5%	5%	7%
Don't know/ not applicable	189	132	87	31	14	25	19	12	31	27	14	13	15	19	8	73	59	28	8	157
	9%	9%	9%	8%	7%	9%	10%	18%	7%	8%	6%	5%	8%	13%	10%	10%	11%	7%	3%	11%
Mean	0.52	0.54	0.49	0.59	0.69	0.48	0.50	0.27	0.26	0.39	0.58	0.58	0.52	0.48	0.21	0.44	0.69	0.61	0.72	0.47
Standard deviation	0.95	0.96	0.98	0.95	0.89	0.85	0.94	0.92	1.00	0.91	0.97	0.92	0.90	1.04	0.80	0.94	0.93	0.92	0.94	0.95
Standard error	0.02	0.03	0.03	0.05	0.07	0.06	0.07	0.12	0.05	0.05	0.06	0.06	0.06	0.09	0.09	0.03	0.05	0.04	0.06	0.03

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 26

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

Having personalised packaging from your pharmacist that shows when each dose should be taken

Base: All respondents

	Gender			Age						Social Grade				Employment Sector	
	Total	Male	Female	18-24	25-34	35-44	45-54	55-64	65+	AB	C1	C2	DE	Public	Private
Unweighted base	2048	1043	1005	202	341	314	379	300	512	581	582	314	571	204	631
Weighted base	2048	997	1051	238	355	336	365	295	459	551	551	446	500	274	885
NET: More likely	1078 53%	529 53%	549 52%	139 58%	186 52%	176 52%	189 52%	141 48%	246 54%	286 52%	281 51%	249 56%	262 52%	155 57%	481 54%
Much more likely	(+2) 453 22%	228 23%	225 21%	49 21%	72 20%	72 22%	77 21%	57 19%	125 27%	99 18%	117 21%	118 26%	119 24%	60 22%	179 20%
Slightly more likely	(+1) 625 31%	301 30%	324 31%	90 38%	114 32%	104 31%	113 31%	84 28%	121 26%	186 34%	164 30%	131 29%	143 29%	95 35%	302 34%
Neither more likely nor less likely	(0) 812 40%	393 39%	419 40%	69 29%	117 33%	138 41%	153 42%	141 48%	194 42%	227 41%	226 41%	162 36%	197 39%	96 35%	330 37%
Slightly less likely	(-1) 27 1%	15 1%	12 1%	8 3%	7 2%	5 1%	4 1%	1 *	2 1%	9 2%	8 1%	2 1%	8 2%	5 2%	15 2%
Much less likely	(-2) 15 1%	11 1%	4 *	- -	4 1%	4 1%	2 1%	3 1%	1 *	2 *	1 *	2 1%	10 2%	2 1%	8 1%
NET: Less likely	42 2%	26 3%	16 2%	8 3%	11 3%	9 3%	6 2%	5 2%	4 1%	11 2%	9 2%	5 1%	18 4%	7 2%	23 3%
Don't know/ not applicable	116 6%	50 5%	66 6%	22 9%	40 11%	13 4%	17 5%	8 3%	15 3%	28 5%	35 6%	30 7%	23 5%	16 6%	51 6%
Mean	0.76	0.76	0.77	0.83	0.77	0.73	0.74	0.67	0.83	0.71	0.75	0.86	0.74	0.80	0.76
Standard deviation	0.85	0.87	0.83	0.82	0.87	0.86	0.83	0.84	0.86	0.80	0.82	0.85	0.92	0.84	0.84
Standard error	0.02	0.03	0.03	0.06	0.05	0.05	0.04	0.05	0.04	0.03	0.04	0.05	0.04	0.06	0.03

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 27

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

Having personalised packaging from your pharmacist that shows when each dose should be taken

Base: All respondents

	Total	Region											
		Scotland	Wales	NET: England	North East	North West	Yorkshire & Humberside	West Midlands	East Midlands	Eastern	London	South East	South West
Unweighted base	2048	174	111	1763	78	244	178	198	154	190	214	311	196
Weighted base	2048	178	104	1766	86	235	174	184	151	198	270	286	180
NET: More likely	1078 53%	109 61%	54 51%	915 52%	44 51%	130 55%	99 57%	104 56%	64 42%	105 53%	138 51%	136 48%	95 53%
Much more likely (+2)	453 22%	50 28%	25 24%	378 21%	25 29%	60 25%	37 21%	47 25%	36 24%	37 19%	53 20%	52 18%	31 17%
Slightly more likely (+1)	625 31%	58 33%	29 28%	538 30%	19 22%	71 30%	62 35%	57 31%	28 18%	68 34%	85 31%	84 29%	64 35%
Neither more likely nor less likely (0)	812 40%	61 34%	46 44%	705 40%	38 45%	87 37%	59 34%	61 33%	76 50%	79 40%	104 38%	133 46%	69 38%
Slightly less likely (-1)	27 1%	3 2%	1 1%	24 1%	2 2%	- -	2 1%	3 1%	5 4%	3 2%	3 1%	2 1%	3 2%
Much less likely (-2)	15 1%	1 1%	1 1%	13 1%	- -	3 1%	1 1%	3 2%	* *	- -	4 2%	2 1%	- -
NET: Less likely	42 2%	4 2%	2 2%	37 2%	2 2%	3 1%	3 2%	6 3%	6 4%	3 2%	8 3%	4 1%	3 2%
Don't know/ not applicable	116 6%	5 3%	3 3%	108 6%	2 2%	16 7%	13 8%	13 7%	6 4%	11 5%	21 8%	13 5%	13 7%
Mean	0.76	0.89	0.75	0.75	0.80	0.84	0.82	0.83	0.65	0.74	0.72	0.67	0.74
Standard deviation	0.85	0.86	0.86	0.85	0.90	0.87	0.83	0.91	0.91	0.79	0.87	0.81	0.78
Standard error	0.02	0.07	0.08	0.02	0.10	0.06	0.06	0.07	0.07	0.06	0.06	0.05	0.06

Medication Adherence Survey

ONLINE Fieldwork: 31st July - 2nd August 2015

Absolutes/col percents

Table 28

Q.7 Which of the following, if any, would impact how likely you would be to adhere to the programme prescribed by your doctor or healthcare professional?

Having personalised packaging from your pharmacist that shows when each dose should be taken

Base: All respondents

	Last Prescribed on NHS								Condition Have Received NHS Prescription For								Medication Adherence			
	Total	NET: In the last year	In the last month	More than 1 month and up to 6 months ago	More than 6 months and up to 1 year ago	More than 1 year and up to 5 years ago	More than 5 years ago	I have never been prescribed any medicine on the NHS	High Blood Pressure	High Cholesterol	Asthma	Mental Health condition	Arthritis	Type 2 Diabetes	Heart Disease	Any other condition	None	Missed a round of medication they are meant to take daily	Not finished a course of medication, such as antibiotics	Never missed a round or not finished a course of medication
Unweighted base	2048	1552	1018	347	187	246	179	71	460	370	244	281	223	171	87	829	445	446	225	1453
Weighted base	2048	1505	925	381	198	279	197	68	417	319	248	247	202	147	81	756	517	431	239	1453
NET: More likely	1078	800	476	207	117	143	110	24	205	159	140	115	113	82	35	400	297	238	121	759
	53%	53%	51%	54%	59%	51%	56%	36%	49%	50%	57%	46%	56%	56%	43%	53%	58%	55%	51%	52%
Much more likely (+2)	453	358	240	84	35	54	33	7	109	84	66	57	56	50	19	175	107	96	48	319
	22%	24%	26%	22%	17%	19%	17%	11%	26%	26%	27%	23%	27%	34%	24%	23%	21%	22%	20%	22%
Slightly more likely (+1)	625	442	236	124	82	89	77	17	96	75	74	57	58	32	16	225	191	141	73	440
	31%	29%	26%	32%	42%	32%	39%	25%	23%	23%	30%	23%	29%	22%	20%	30%	37%	33%	31%	30%
Neither more likely nor less likely (0)	812	599	382	150	67	116	70	27	183	148	89	118	83	54	41	306	169	173	101	571
	40%	40%	41%	39%	34%	42%	36%	40%	44%	46%	36%	48%	41%	37%	50%	40%	33%	40%	42%	39%
Slightly less likely (-1)	27	16	8	6	2	-	6	6	6	2	6	3	1	-	-	8	4	3	6	18
	1%	1%	1%	1%	1%	-	3%	9%	1%	*	2%	1%	*	-	-	1%	1%	1%	2%	1%
Much less likely (-2)	15	15	14	1	-	-	-	-	12	2	6	-	2	2	4	-	3	4	7	*
	1%	1%	1%	*	-	-	-	-	3%	1%	2%	-	1%	1%	*	-	1%	2%	2%	*
NET: Less likely	42	31	22	7	2	-	6	6	17	4	11	3	3	2	-	12	4	6	10	26
	2%	2%	2%	2%	1%	-	3%	9%	4%	1%	5%	1%	1%	1%	-	2%	1%	2%	4%	2%
Don't know/ not applicable	116	75	45	17	12	19	11	11	13	8	7	12	3	9	5	38	46	14	7	97
	6%	5%	5%	5%	6%	7%	5%	16%	3%	3%	3%	5%	1%	6%	6%	5%	9%	3%	3%	7%
Mean	0.76	0.78	0.77	0.78	0.81	0.76	0.74	0.45	0.70	0.76	0.79	0.71	0.82	0.94	0.71	0.78	0.85	0.78	0.67	0.77
Standard deviation	0.85	0.87	0.91	0.83	0.74	0.78	0.78	0.86	0.97	0.88	0.96	0.85	0.88	0.93	0.85	0.84	0.77	0.84	0.90	0.84
Standard error	0.02	0.02	0.03	0.05	0.06	0.05	0.06	0.11	0.05	0.05	0.06	0.05	0.06	0.07	0.09	0.03	0.04	0.04	0.06	0.02